

# Mitogen requirement for cell cycle progression in the absence of pocket protein activity

Floris Fojjer,<sup>1</sup> Rob M.F. Wolthuis,<sup>1</sup> Valerie Doodeman,<sup>1</sup> René H. Medema,<sup>2</sup> and Hein te Riele<sup>1,\*</sup>

<sup>1</sup> Division of Molecular Biology, The Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands

<sup>2</sup> Present address: Experimental Oncology, University Medical Center, Stratenum 2.103, Universiteitsweg 100, 3584 CG Utrecht, The Netherlands

\*Correspondence: [h.t.rielle@nki.nl](mailto:h.t.rielle@nki.nl)

## Summary

**Primary mouse embryonic fibroblasts lacking expression of all three retinoblastoma protein family members (TKO MEFs) have lost the G<sub>1</sub> restriction point. However, in the absence of mitogens these cells become highly sensitive to apoptosis. Here, we show that TKO MEFs that survive serum depletion pass G<sub>1</sub> but completely arrest in G<sub>2</sub>. p21<sup>CIP1</sup> and p27<sup>KIP1</sup> inhibit Cyclin A-Cdk2 activity and sequester Cyclin B1-Cdk1 in inactive complexes in the nucleus. This response is alleviated by mitogen restimulation or inactivation of p53. Thus, our results disclose a cell cycle arrest mechanism in G<sub>2</sub> that restricts the proliferative capacity of mitogen-deprived cells that have lost the G<sub>1</sub> restriction point. The involvement of p53 provides a rationale for the synergism between loss of *Rb* and *p53* in tumorigenesis.**

## Introduction


Proliferation of cells in culture is dependent on the presence of mitogenic stimuli. In the absence of mitogens, cells fail to progress through the so-called restriction point and arrest in G<sub>0</sub>/G<sub>1</sub>. The restriction point was originally defined by Pardee as the event in G<sub>1</sub> after which cells are committed to progress through the cell cycle independently of mitogens (Pardee, 1974). Most physiological signals that force cells to pass the G<sub>1</sub> restriction point activate Ras signaling, ultimately resulting in increased transcription of *Cyclin D1*. Additionally, Cyclin D1 protein levels are stabilized by activation of the PI3 kinase pathway (Malumbres and Barbacid, 2001; Schmidt et al., 2002). Cyclin D1 with its catalytic subunits, the cyclin-dependent kinases (Cdks) Cdk4 and Cdk6, mediates the extensive phosphorylation and partial inactivation of the retinoblastoma protein pRb during G<sub>1</sub> (Planas-Silva and Weinberg, 1997). pRb hyperphosphorylation triggers the activity of E2F transcription factors, leading to enhanced Cyclin E levels. Increased Cyclin E/Cdk2 activity allows further pRb hyperphosphorylation, G<sub>1</sub> progression, and initiation of S phase. Conversely, mitogen starvation results in reduced transcription of *Cyclin D1*, destabilization of Cyclin D1 protein, and induction of the Cdk2 inhibitor p27<sup>KIP1</sup>, resulting in hypophosphorylated pRb, subsequent inhibition of E2F activity, and arrest in G<sub>0</sub>/G<sub>1</sub> (Blagosklonny and Pardee, 2002).

Despite the critical role of pRb in controlling the G<sub>1</sub> restriction point, pRb knockout primary mouse embryonic fibroblasts (MEFs) do not proliferate in the absence of mitogens but still arrest (Herrera et al., 1996; Almasan et al., 1995). This could be explained by the redundant functions of the pRb family members p107 and p130 (Mulligan and Jacks, 1998), since these so-called pocket proteins collectively regulate the activity of E2F transcription factors. We and others have shown that complete disruption of the pocket proteins in MEFs (*triple* knockout MEFs or TKO MEFs) resulted in abrogation of the G<sub>1</sub> restriction point (Dannenberg et al., 2000; Sage et al., 2000). Also, viral oncoproteins like E1A, E7, or Large T, which are encoded by adenoviruses, human papillomaviruses, and polyomaviruses, respectively, can overcome the G<sub>1</sub> restriction point by direct interaction with the pRb family members (Helt and Galloway, 2003).

The function of pocket proteins is often compromised in human cancer through either expression of viral oncogenes, overexpression of *Cyclin D1*, or loss of the tumor suppressor gene *p16<sup>INK4A</sup>*, suggesting that loss of the G<sub>1</sub> restriction point is a critical step in the development of cancer (Pardee, 1974; Malumbres and Barbacid, 2001). However, here we show that ablation of the pocket proteins in primary cells in vitro does not result in complete alleviation of cell cycle control. We provide evidence for a p53-dependent control mechanism that acts in the G<sub>2</sub>

## SIGNIFICANCE

**Cancer is a multistep process resulting from abrogation of several barriers to uncontrolled proliferation. One of these barriers is formed by the retinoblastoma protein family that prevents cells from undesired replication of DNA. We have identified a cell cycle barrier that prevents unbalanced cell division of mitogen-starved cells that have passed the restriction point. We show that loss of these barriers, e.g., by combined inactivation of retinoblastoma proteins and p53, allows proliferation of cells in the absence of growth stimuli. This observation gives insight into the causes of oncogenic transformation, providing opportunities for therapeutic intervention.**



**Figure 1.** Serum starvation of TKO MEFs results in massive apoptosis from  $G_2$  and a  $G_2$  arrest in the surviving cells

**A:** Apoptosis induction in wild-type and TKO MEFs upon serum starvation. Graphs represent the average of three independent experiments; error bars represent standard deviations.

**B:** Cell cycle distribution of serum-starved wild-type MEFs. Graphs represent the average of three independent experiments; error bars represent standard deviations.

**C and D:** Induction of apoptosis in asynchronously (**C**) and synchronously (**D**) growing TKO MEFs by serum depletion. Time is given in hh:mm after serum starvation.

**E:** S phase progression of TKO MEFs in the presence and absence of serum after mitosis as determined by BrdU incorporation.

**F and G:** Cell cycle distribution of serum-starved TKO (**F**) and TKO-Bcl2 MEFs (**G**). Graphs represent the average of three independent experiments; error bars represent standard deviations.


phase of the cell cycle to arrest or eliminate serum-deprived cells lacking a functional  $G_1$  restriction point.

## Results

### Mitogen starvation in the absence of pocket proteins results in massive apoptosis and arrest in $G_2$

Consistent with previous results (Sage et al., 2000; Dannenberg et al., 2000), serum deprivation in wild-type MEFs led to the accumulation of cells with a  $2n$  and, to a lesser extent,  $4n$  DNA content, but not to apoptosis (Figures 1A and 1B). In contrast, TKO MEFs underwent massive apoptosis upon serum starvation (Figure 1A). We used time-lapse microscopy to monitor the onset of apoptosis in serum-starved TKO MEF cultures. Several

hours after serum withdrawal, the first apoptotic events could be observed (floating cells, Figure 1C). To examine if programmed cell death occurred at a specific point in the cell cycle, we synchronized TKO MEFs by mitotic shake-off and cultured them under serum-free or high-serum conditions. Time-lapse microscopy revealed that, in the absence of serum, the first apoptotic events (indicated by arrows in Figure 1D) occurred at approximately 12 hr after exit from mitosis. Simultaneously, we followed the progression of synchronized cells by measuring 5-bromodeoxyuridine (BrdU) incorporation. Figure 1E shows that the kinetics of progression through  $G_1$  and entry into S phase of TKO cells were independent of serum, consistent with loss of the  $G_1$  restriction point (Dannenberg et al., 2000; Sage et al., 2000). The appearance of apoptotic cells coincided with



**Figure 2.** Serum starvation-induced  $G_2$  arrest in apoptosis-resistant TKO MEFs is associated with  $p21^{CIP1}$ - and  $p27^{KIP1}$ -dependent inhibition of Cyclin A and B1 kinase activity

**A:** Cyclin A- and B1-associated kinase activities upon serum starvation and total levels of cyclins and Cdk1. Graphs represent the average values of three independent experiments, and error bars represent standard deviations; blots represent the results from a representative experiment.

**B:** Total levels of  $p21^{CIP1}$  and  $p27^{KIP1}$  in serum-starved apoptosis-resistant TKO MEFs. Cdk4 functions as a loading control.

**C–E:** Immunoprecipitations showing interactions of  $p21^{CIP1}$  and  $p27^{KIP1}$  with Cyclin B1 (**C** and **D**) and Cyclin A (**E**) from serum-starved TKO-Bcl2 MEFs.

**F:** Total Cdk2 levels and Cdk2 kinase activity in serum-starved TKO-Bcl2 MEFs. Percentages represent quantified values of labeled Histone H1.

**G:** Immunoprecipitation of Cdk2 from serum-starved TKO-Bcl2 MEFs.

exit from S phase and onset of  $G_2$ . Clearly, this apoptotic response could act as an effective safety strategy to prevent propagation of  $G_1$  restriction point-defective cells in the absence of mitogens. Strikingly, however, we observed that the small fraction of TKO MEFs that survived serum starvation did not proliferate but arrested in the  $G_2$  phase of the cell cycle (Figure 1F). In order to study this cell cycle arrest in detail, we generated apoptosis-resistant TKO MEFs by expressing the antiapoptotic protein Bcl2 using retroviral transduction. TKO-Bcl2 cells largely survived serum starvation but arrested in  $G_2$  with the same kinetics as TKO MEFs (Figure 1G).


#### Inhibition of Cyclin A- and B1-associated kinase activity

Cdk1 kinase activity is essential for progression into mitosis and is dependent on association with Cyclin A or Cyclin B1 (Nigg, 2001). We therefore determined these Cyclin-associated kinase activities in apoptosis-resistant TKO MEFs after serum starvation. Indeed, Cyclin B1-Cdk1 kinase activity was downmodulated 5-fold compared to B1-associated kinase activity in asynchronously growing cultures, although total levels of Cyclin B1 and Cdk1 remained constant. Also, Cyclin A-associated kinase activity was reduced upon serum starvation, while the levels of Cyclin A slightly increased (Figure 2A). Cyclin-associated kinases can be inhibited by cyclin-dependent kinase inhibitors (CKIs) (Coqueret, 2003). We therefore determined the levels of the CKIs  $p21^{CIP1}$  and  $p27^{KIP1}$  in serum-starved apoptosis-resistant TKO MEFs. Figure 2B shows that both  $p21^{CIP1}$  and  $p27^{KIP1}$  levels were strongly increased upon serum starvation. Previous reports have suggested that  $p21^{CIP1}$  can inhibit Cyclin B1-associated kinase activity by direct interaction (Smits et al., 2000b; Baus et al., 2003). Figure 2C shows that upon

serum starvation the amount of Cyclin B1 coprecipitating with  $p21^{CIP1}$  increased and reached a maximum level after 5 and 7 days of serum starvation. Similarly,  $p21^{CIP1}$  increasingly coimmunoprecipitated with Cyclin B1-Cdk1. Thus, Cyclin B1-Cdk1 and  $p21^{CIP1}$  were present in the same complex at 5 and 7 days after serum deprivation (Figure 2D). Interestingly, we observed that also  $p27^{KIP1}$  was sequestered into Cyclin B1 complexes in these cells, although here the level of interaction reached a maximum after 3–5 days of serum starvation and then decreased (Figure 2D, lower panel). Since  $p27^{KIP1}$  was absent in  $p21^{CIP1}$  immunoprecipitates, we hypothesize that  $p21^{CIP1}$  and  $p27^{KIP1}$  compete for similar Cyclin B1-Cdk1 binding sites (data not shown). In parallel, we found that also Cyclin A increasingly associated with  $p21^{CIP1}$  and  $p27^{KIP1}$  upon serum withdrawal (Figure 2E). In agreement with this, increasing levels of  $p21^{CIP1}$  and  $p27^{KIP1}$  also inhibited the kinase activity of Cdk2 (Figure 2F) through direct association (Figure 2G). Thus, inhibition of Cyclin A-Cdk2 may contribute to the  $G_2$  arrest, which is consistent with a role for Cyclin A-Cdk2 in progression from  $G_2$  to mitosis (Furuno et al., 1999).

#### Individual roles of the CKIs in the $G_2$ arrest

To evaluate the individual roles of  $p21^{CIP1}$  and  $p27^{KIP1}$  in the  $G_2$  arrest, we asked whether high levels of  $p27^{KIP1}$  could block TKO-Bcl2 MEFs in  $G_2$ . We synchronized TKO-Bcl2 MEFs by mitotic shake-off and microinjected YFP-tagged  $p27^{KIP1}$  12 hr later (Connor et al., 2003), when the majority of the cells were in  $G_2$ . Time-lapse microscopy showed that control-injected  $G_2$  cells readily progressed through mitosis, whereas none of the YFP- $p27^{KIP1}$ -expressing  $G_2$  cells entered mitosis, indicating that  $p27^{KIP1}$  is capable of establishing a  $G_2$  arrest (data not shown).



**Figure 3.** Inhibition of p21<sup>CIP1</sup> delays the G<sub>2</sub> arrest **A:** Total levels of Cyclin B1, p27<sup>KIP1</sup>, and p21<sup>CIP1</sup> in pRS control- and pRS-p21<sup>CIP1</sup>-infected TKO-Bcl2 MEFs upon serum starvation. Actin functions as a loading control.

**B and C:** Immunoprecipitations of Cyclin B1 (**B**) and Cdk2 (**C**) from serum-starved control and pRS-p21<sup>CIP1</sup>-infected TKO Bcl2 MEFs probed for the presence of p21<sup>CIP1</sup> and p27<sup>KIP1</sup>.

**D and E:** Cyclin B1-associated kinase activity (**D**) and Cdk2 kinase activity (**E**) in serum-starved TKO-Bcl2 MEFs infected with either pRS control or pRS-p21<sup>CIP1</sup> viruses. Error bars represent standard deviations of two independent experiments.

**F:** G<sub>2</sub> accumulation of serum-starved pRS control- or pRS-p21<sup>CIP1</sup>-infected TKO-Bc2 MEFs as determined by FACS analysis. Graphs represent the average of two independent experiments; error bars represent standard deviations.

In another approach, we have attempted to knock down the CKIs using vector-driven RNAi constructs. Although we were unsuccessful in knocking down p27<sup>KIP1</sup> in TKO-Bcl2 MEFs, we succeeded in knocking down p21<sup>CIP1</sup>. Figure 3A shows that p21<sup>CIP1</sup> was hardly induced in TKO-Bcl2 MEFs expressing pRetroSuper p21<sup>CIP1</sup> upon serum starvation, whereas it readily accumulated in control-infected cells. pRetroSuper p21<sup>CIP1</sup> did not affect Cyclin B1 expression and allowed normal p27<sup>KIP1</sup> accumulation upon serum starvation (Figure 3A). Inhibition of p21<sup>CIP1</sup> expression resulted in less p21<sup>CIP1</sup> associated to Cyclin B1 (Figure 3B, lower panel) and Cdk2 (Figure 3C, lower panel). This resulted in a strong increase of Cyclin B1-associated and Cdk2 kinase activities (Figures 3D and 3E) in cells growing in high serum. During serum starvation, these activities were reduced, most likely due to the strong association of p27<sup>KIP1</sup> to Cyclin B1 and Cdk2 (Figures 3B and 3C), but not to the same level as in pRS control cells (Figures 3D and 3E). Consistently, FACS analysis revealed that the accumulation of cells in G<sub>2</sub> was reduced upon expression of p21<sup>CIP1</sup>-RNAi (Figure 3F). As cell numbers only moderately increased (data not shown), these data suggest that inhibition of p21<sup>CIP1</sup> delayed the G<sub>2</sub> arrest but was not sufficient for unconstrained proliferation under mitogen-deprived conditions.

These results demonstrate that in serum-starved TKO MEFs association of both p21<sup>CIP1</sup> and p27<sup>KIP1</sup> to Cyclin B1-Cdk1 complexes and Cdk2 contribute to inhibition of mitotic cyclins, leading to a sustained G<sub>2</sub> arrest.


#### The mitogen starvation-induced G<sub>2</sub> arrest is reversible

If the G<sub>2</sub> arrest displays typical characteristics of a restriction point, serum-starved apoptosis-resistant TKO MEFs should be

capable of reentering the cell cycle upon serum addition. Time-lapse microscopy revealed the appearance of mitotic cells 18–24 hr after serum readdition to G<sub>2</sub>-arrested cells (data not shown). Using MPM2 as a marker for the appearance of mitotic cells, we confirmed by FACS that the cells entered mitosis at 18–24 hr after serum restimulation (Figure 4A). To verify the relevance of complexes of Cyclin B1 with p21<sup>CIP1</sup> and p27<sup>KIP1</sup>, we determined whether they were affected by serum readdition. p27<sup>KIP1</sup> completely disappeared from the Cyclin B1 kinase complex within 15 hr after the addition of serum (Figure 4B). p21<sup>CIP1</sup>-Cyclin B1 complexes slightly increased after 15 hr but gradually decreased when the cells entered mitosis, although they did not completely disappear. Consistently, p27<sup>KIP1</sup> levels dropped promptly, whereas p21<sup>CIP1</sup> levels only moderately decreased. Importantly, Cyclin A-associated kinase activity reappeared within 15 hr after serum addition, reaching a maximum at 21 hr, and Cyclin B1-Cdk1 was reactivated within 18 hr and peaked at 24 hr after serum stimulation (Figure 4C). These results show that mitogen stimulation results in release of CKIs, reactivation of Cyclin A- and Cyclin B1-associated kinase activities, G<sub>2</sub> progression, and mitotic entry, demonstrating that the G<sub>2</sub> arrest is reversible.

#### The mitogen starvation-induced G<sub>2</sub> arrest is different from a DNA damage response

During G<sub>2</sub> phase and during a DNA damage-induced G<sub>2</sub> arrest, Cyclin B1 localizes to the cytoplasmic compartment in an inactive state (Toyoshima et al., 1998). We wanted to compare the G<sub>2</sub> serum starvation arrest with a DNA damage-induced arrest. In TKO MEFs, DNA damage indeed induced cytoplasmic retention of Cyclin B1. In contrast, serum starvation resulted in



**Figure 4.** The serum starvation-induced  $G_2$  arrest is reversible

**A:** Mitotic entry measured by MPM2 positivity of serum-starved TKO-Bcl2 MEFs upon serum readdition. Graph represents average values of two independent experiments; error bars show standard deviations.

**B:** Disruption of the CKI/Cyclin B1 kinase complexes in serum-stimulated TKO-Bcl2 MEFs. Lower panel shows whole-cell lysates. Cdk4 functions as a loading control.

**C:** Induction of Cyclin A- and B1-associated kinase activities upon serum stimulation of  $G_2$ -arrested/apoptosis-resistant TKO MEFs. Graph represents average values of two independent experiments; error bars show standard deviations.

accumulation of Cyclin B1 in the nucleus (Figure 5A). Additionally, inhibition of the DNA damage-induced kinases ATM/ATR by caffeine could not release the cells from the  $G_2$  block, whereas serum readdition resulted in progression into mitosis (Figure 5B). Although inhibition of ATM/ATR kinase activity did not result in mitotic entry, we found that p21<sup>CIP1</sup> almost completely disappeared from the Cyclin B1-Cdk1 kinase complex (Figure 5C, upper panel) and that total p21<sup>CIP1</sup> levels moderately decreased (Figure 5C, lower panel). However, total levels of p27<sup>KIP1</sup> as well as the amount of p27<sup>KIP1</sup> associated to Cyclin B1 increased in caffeine-treated cells, in contrast to serum-stimulated cells, where p27<sup>KIP1</sup> levels and Cyclin B1-associated p27<sup>KIP1</sup> levels dramatically dropped (Figure 5C). Concordantly, we found that inhibition of ATM/ATR kinase activities was not sufficient to reactivate Cyclin B1- and Cyclin A-associated kinase activities under mitogen-deprived conditions (Figure 5D). This suggests that p21<sup>CIP1</sup> but not p27<sup>KIP1</sup> is induced by ATM/ATR-dependent signaling upon serum starvation in apoptosis-resistant TKO MEFs. Therefore, we hypothesize that the serum starvation-induced  $G_2$  arrest might partially depend on the DNA damage signaling pathway for the induction of p21<sup>CIP1</sup>. However, when the DNA damage pathway is inhibited during an already established  $G_2$  arrest, p27<sup>KIP1</sup> is sufficient to prevent reactivation of Cyclin B1- and Cyclin A-associated kinases and alleviation of the  $G_2$  arrest.


#### Increased inhibition of pocket protein activity correlates with increased $G_2$ arrest

To determine whether the  $G_2$  arrest also occurs in MEFs lacking one or two pocket proteins, we compared the serum starvation responses of wild-type,  $Rb^{-/-}$ ,  $p107^{-/-}p130^{-/-}$ ,  $Rb^{-/-}p130^{-/-}$ ,

$Rb^{-/-}p107^{-/-}$ , and apoptosis-resistant TKO MEFs by determining the  $G_2$  over  $G_1$  ratios. While  $p107^{-/-}p130^{-/-}$  MEFs responded in the same way as wild-type MEFs, loss of  $Rb$  alone resulted in an increased  $G_2/G_1$  ratio (0.69 versus 0.45 in wild-type MEFs) (Figure 6A, upper three histograms).  $Rb^{-/-}p130^{-/-}$  MEFs behaved like  $Rb^{-/-}$  MEFs, whereas combinational loss of  $p107$  and  $Rb$  resulted in an increased  $G_2/G_1$  ratio when serum-starved (1.1 versus 0.69 in  $Rb$ -deficient MEFs; Figure 6A, lower three histograms). Serum-starved  $Rb^{-/-}p107^{-/-}$  MEFs showed a substantial fraction of polyploid cells, whereas this feature was less prominent in apoptosis-resistant TKO MEFs. Therefore, a part of the 4n cells observed in the serum-starved  $Rb^{-/-}p107^{-/-}$  MEFs may be tetraploid cells that have arrested in  $G_1$ , while the 8n cell population contributes to the  $G_2$  fraction. We did not observe a substantial induction of apoptosis in either of the cell cultures (indicated by "A" in Figure 6A).

We next determined the complex formation between Cyclin B1 and p27<sup>KIP1</sup> and Cyclin B1 and p21<sup>CIP1</sup> in serum-starved cells. The p27<sup>KIP1</sup>-Cyclin B1 interaction could be detected in all serum-starved knockout MEFs except for the  $p107^{-/-}p130^{-/-}$  MEFs (Figure 6B) and correlated with the fraction of  $G_2$  cells. Since no Cyclin B1 could be detected in serum-starved wild-type and  $p107^{-/-}p130^{-/-}$  MEFs (Figure 6B), we envisage that the cells arrested with a 4n DNA content are tetraploid cells arrested in  $G_1$ , a described tendency for MEFs upon passaging (Borel et al., 2002).

These results suggest that also in MEFs in which the pocket protein family is only partially ablated, serum starvation induces a  $G_2$  arrest that is mediated by a p27<sup>KIP1</sup>-Cyclin B1 interaction. However, the serum starvation-induced association of p21<sup>CIP1</sup> to Cyclin B1 appears to be unique for TKO MEFs (Figure 6B).



**Figure 5.** The G<sub>2</sub> arrest is partially different from a DNA damage response

**A:** Double immunofluorescence staining for Cyclin B1 (green) and DNA (blue) showing DNA damage-induced cytoplasmic localization and serum starvation-induced nuclear localization of Cyclin B1.

**B:** Mitotic entry determined by MPM2 positivity of serum-starved TKO-Bcl2 MEFs stimulated with either 10% serum (white bars) or 5 mM caffeine (black bars). Graph represents average values of two independent experiments; error bars show standard deviations.

**C:** p21<sup>CIP1</sup>-Cyclin B1 and p27<sup>KIP1</sup>-Cyclin B1 complex formation in serum-starved TKO-Bcl2 MEFs stimulated with either 10% serum or 5 mM caffeine. Cdk1 functions as a loading control.

**D:** Cyclin A- and B1-associated kinase activities of serum-starved TKO-Bcl2 MEFs treated with 5 mM caffeine. Values represent relative kinase activities as compared to asynchronously proliferating TKO-Bcl2 MEFs. Error bars show standard deviations of two experiments.

In human cancer, the G<sub>1</sub> restriction point is often affected by mechanisms other than genetic ablation of pocket proteins, such as expression of viral oncogenes, loss of p16<sup>INK4A</sup>, or mutation or amplification of *Cyclin D1*. Therefore, we investigated whether the responses to serum starvation of p16<sup>INK4A</sup>-/- MEFs, wild-type MEFs expressing a nondegradable form of Cyclin D1 (Agami and Bernards, 2000), and wild-type MEFs stably expressing E1A were similar to that of pocket protein knockout MEFs. Since E1A-expressing MEFs underwent massive apoptosis as previously described (data not shown) (Mymryk et al., 1994), we introduced Bcl2 in these cells. p16<sup>INK4A</sup> null MEFs responded like wild-type MEFs upon serum starvation: they arrested in G<sub>1</sub> phase of the cell cycle and had completely lost Cyclin B1 expression (data not shown). In contrast, MEFs expressing E1A and Bcl2 accumulated in G<sub>2</sub> almost to the same extent as apoptosis-resistant TKO MEFs (Figure 6C). This arrest correlated with inhibition of Cyclin B1-associated kinase activity (Figure 6E) and binding of p27<sup>KIP1</sup> and p21<sup>CIP1</sup> to Cyclin B1 (Figure 6F). Although in E1A-expressing cells a clear interaction of p21<sup>CIP1</sup> and Cyclin B1 was already present in proliferating cells and the levels of Cyclin B1 decreased during serum starvation, the fraction of Cyclin B1 associated to p21<sup>CIP1</sup> increased upon mitogen-starved conditions (Figure 6F).


MEFs expressing a nondegradable form of Cyclin D1 accumulated in G<sub>2</sub> to the same extent as *Rb*<sup>-/-</sup> MEFs, and approximately 10% of the cells underwent apoptosis when serum starved (Figure 6D). G<sub>2</sub> arrest correlated with loss of Cyclin B1

kinase activity (Figure 6E) through binding of the CKIs (Figure 6F).

Together, these results show that interference with the G<sub>1</sub> restriction point by events frequently seen in tumor development is not sufficient for unhindered proliferation under mitogen-deprived conditions. Instead, such cells arrest in G<sub>2</sub> due to inhibition of Cyclin B1 by p21<sup>CIP1</sup> and/or p27<sup>KIP1</sup>.

### p53 is essential for the G<sub>2</sub> arrest

p53 is an important mediator of apoptosis and growth arrest under many stress conditions (Hanahan and Weinberg, 2000). To investigate the role of p53 in apoptosis and growth arrest of serum-deprived TKO MEFs, we inactivated p53 function in TKO MEFs by RNA interference using pRetroSuper-p53. Figure 7A shows that, in TKO-p53RNAi MEFs, the levels of p53 and one of its target proteins, p21<sup>CIP1</sup>, were strongly reduced compared to the levels in control-infected TKO and apoptosis-resistant TKO-Bcl2 MEFs. Knocking down p53 expression almost completely prevented apoptosis in serum-starved TKO MEFs and was even more effective as the expression of Bcl2 (Figure 7B). Furthermore, unlike TKO-Bcl2 MEFs, serum-starved TKO-p53RNAi MEFs continued to incorporate BrdU while the accumulation of cells in G<sub>2</sub> was strongly diminished (Figure 7C). Importantly, simultaneous ablation of pocket proteins and p53 resulted in prolonged proliferation in the absence of mitogens (Figure 7D; Figure S1 in the Supplemental Data available with this article online): TKO-p53RNAi cells grew



**Figure 6.** Increasing pocket protein loss results in increased G<sub>2</sub> arrest and increased p27<sup>KIP1</sup>-Cyclin B1 complex formation under serum-starved conditions. **A:** Histograms showing cell cycle distribution of pocket protein knockout MEFs at 3 days of serum starvation ("A" refers to the apoptotic population [ $<2n$ ]). **B:** Immunoprecipitation of Cyclin B1 in serum-starved wild-type and pocket protein knockout MEFs. Actin in whole-cell lysates functions as a loading control. **C and D:** Histograms showing cell cycle distributions of wild-type MEFs expressing either E1A and Bcl2 (**C**) or a nondegradable form of Cyclin D1 (Cyclin D1<sup>ND</sup>) (**D**). **E:** Cyclin B1-associated kinase activity of serum-starved E1A-Bcl2 MEFs and Cyclin D1<sup>ND</sup> MEFs. Percentages represent a quantification of the intensity of the kinase activity. **F:** Immunoprecipitations (upper panels) and whole-cell lysates (lower panels) from Cyclin B1 of serum-starved wild-type, E1A-Bcl2, and Cyclin D1<sup>ND</sup> MEFs. Actin functions as a loading control.


exponentially for at least three population doublings in the absence of serum, although the generation time had increased from approximately 1 day under high-serum conditions to 2 days. However, when reaching confluency, many TKO-p53 RNAi cells died. Unfortunately, we could not assess the growth potential of TKO-p53RNAi cells for a longer time, as we could not passage these cells under serum-free conditions. Time-lapse microscopy revealed that TKO-p53RNAi MEFs being passaged under serum-free conditions did not reattach and subsequently died, suggesting that serum contains factors essential for cell attachment (data not shown). Alternatively, cell death could be a consequence of residual trypsin activity, which is normally inhibited by serum.

As neither Cyclin A- and B1-associated kinase activities (Figure 7E) nor Cdk2 kinase activity (Figure 7F) in these cells was inhibited until day 7, we compared the presence of p21<sup>CIP1</sup> and p27<sup>KIP1</sup> in the Cyclin B1-Cdk1 complexes in TKO-Bcl2 and TKO-p53RNAi cells. Figure 7G shows that p21<sup>CIP1</sup> and p27<sup>KIP1</sup> were virtually absent from Cyclin B1-Cdk1 kinase complexes isolated

from TKO-p53 RNAi MEFs. Similarly, Cdk2 immunoprecipitates showed no association to p21<sup>CIP1</sup> and a weak interaction to p27<sup>KIP1</sup>, which was not induced by serum starvation (Figure 7H). As is shown in Figure 7I (middle panel), p21<sup>CIP1</sup> expression was almost completely abolished in mitogen-starved TKO-p53RNAi MEFs, while p27<sup>KIP1</sup> expression was markedly reduced (Figure 7I, upper panel). This suggests that in the absence of p53 the accumulation of both p21<sup>CIP1</sup> and p27<sup>KIP1</sup> was largely suppressed, allowing mitogen-independent proliferation of pocket protein-deficient cells.

In another approach, we expressed E1A in p53<sup>-/-</sup> MEFs. Whereas in control-infected cells proliferation was inhibited as assayed by BrdU incorporation, E1A-expressing p53<sup>-/-</sup> MEFs continued to incorporate BrdU up to 7 days of mitogen starvation, although the percentage of BrdU-positive cells gradually decreased (Figure 7J).

In conclusion, these results indicate that the combined abrogation of p53 and pocket protein functions strongly potentiates the ability of cells to proliferate independently of mitogens.



**Figure 7.** Serum starvation-induced apoptosis and  $G_2$  arrest in TKO MEFs are dependent on p53

**A:** Levels of p53 and p21<sup>CIP1</sup> in TKO, TKO-Bcl2, and TKO-p53RNAi MEFs. Cdk1 functions as a loading control.

**B:** Induction of apoptosis by serum starvation in TKO, TKO-p53RNAi, and TKO-Bcl2 MEFs. Values represent the average of two experiments; error bars represent standard deviations.

**C:** Cell cycle distribution of serum-starved TKO-p53RNAi MEFs. Values represent the average of three independent experiments; error bars show standard deviations.

**D:** Growth curves showing serum-independent proliferation of p53-deficient TKO MEFs. Experiment was performed in triplo; error bars show standard deviations.

**E:** Cyclin A- and B1-associated kinase activities upon serum starvation of TKO-p53 RNAi MEFs. Graph represents average values of four independent experiments; error bars show standard deviations.

**F:** Cdk2 kinase activity of serum-starved TKO-p53-RNAi MEFs. Percentages represent quantification of labeled Histone H1.

**G** and **H:** Immunoprecipitations of Cyclin B1 (**G**) and Cdk2 (**H**) from serum-starved TKO-Bcl2 MEFs and TKO-p53RNAi MEFs.

**I:** Whole-cell lysates of serum-starved TKO-Bcl2 and TKO-p53RNAi MEFs. Cdk4 functions as a loading control.

**J:** Proliferating cells as determined by BrdU incorporation in control- and E1A-infected p53<sup>-/-</sup> MEF cultures. Error bars represent standard deviations of two experiments.

## Discussion

Loss of pocket protein function results in deregulated E2F activity and consequently elevated E2F target gene expression. Amongst E2F targets are many proapoptotic genes like Apaf-1, caspases, Bad, and Bak1 (Stanelle et al., 2002; Muller et al., 2001). Serum starvation activates the FOXO family of transcription factors and thereby expression of proapoptotic genes like FasL and Bim (Burgering and Kops, 2002). It is likely that the combinatorial effects of elevated expression of proapoptotic E2F and FOXO target genes cause the massive apoptosis that we have observed in serum-starved TKO MEFs. We have also shown that loss of p53 suppresses serum starvation-induced

apoptosis. A simultaneous requirement for p53 and free E2F as was shown for Apaf-1 induction could explain this observation (Moroni et al., 2001). Our results are consistent with previous reports showing that mitogen-starved E1A-expressing murine cells are highly sensitive to apoptosis (Mymryk et al., 1994), which could be suppressed by inactivation of p53 by adenoviral E1B (Rao et al., 1992).


Protection from apoptosis is a hallmark of cancer. The antiapoptotic protein Bcl2 can suppress apoptosis induced in mitogen-starved fibroblasts that express high levels of the oncogenes c-Myc (Hanahan and Weinberg, 2000) or E1A (Rao et al., 1992). We therefore overexpressed Bcl2 in TKO MEFs and studied the cell cycle progression of these cells upon

mitogen starvation. Indeed, apoptosis was suppressed, but interestingly the majority of surviving cells arrested with a G<sub>2</sub> DNA content. Additionally, we have shown that partial abrogation of the G<sub>1</sub> restriction point by loss of Rb alone, combined loss of Rb and p107, expression of the adenoviral oncogene E1A, or expression of a nondegradable form of Cyclin D1 also resulted in an accumulation of cells in G<sub>2</sub> under mitogen-deprived conditions. On the other hand, loss of the Cyclin D1 inhibitor *p16<sup>INK4A</sup>* did not result in a partially defective G<sub>1</sub> restriction point, which is not surprising, since Cyclin D1 completely disappears under mitogen-starved conditions (Blagosklonny and Pardee, 2002).

Our results show that alleviation of both the G<sub>1</sub> restriction point and the apoptotic response is not sufficient for mitogen-independent proliferation. Instead, cells lacking a functional G<sub>1</sub> restriction point can activate a pocket protein-independent arrest in G<sub>2</sub> that serves as an emergency brake to prevent unconstrained proliferation under mitogen-deprived conditions. We are currently analyzing Rb-deficient premalignant lesions to investigate whether this mechanism operates in vivo.

Serum starvation also induces FOXO-dependent transcription of *p27<sup>KIP1</sup>* (Medema et al., 2000). In wild-type cells, *p27<sup>KIP1</sup>* induction inhibits Cyclin E-Cdk2 activity, which contributes to hypophosphorylation of the pocket proteins and G<sub>1</sub> arrest. Here, we show that in pocket protein-compromised MEFs, induction of *p27<sup>KIP1</sup>* and *p21<sup>CIP1</sup>* upon serum starvation is responsible for the G<sub>2</sub> arrest through inhibitory interactions with Cyclin B1 and Cyclin A. As Cdk2 is not essential for cell cycle progression (Ortega et al., 2003), one may argue that inhibition of Cdk1, rather than Cdk2 is critical for serum starvation-induced G<sub>2</sub> arrest. However, others have shown that Cyclin A-Cdk2 and Cyclin B1-Cdk1 kinase activities are essential for mitotic entry (Furuno et al., 1999; Nurse, 1990; Nigg, 2001). Therefore, we think that inhibition of both activities is relevant to the serum starvation-induced G<sub>2</sub> arrest. This is consistent with our finding that readdition of serum caused reappearance of Cyclin B1- and Cyclin A-associated kinase activities and propagated entry of mitogen-starved cells into mitosis. We also consider the activity of both *p27<sup>KIP1</sup>* and *p21<sup>CIP1</sup>* to be critical to the G<sub>2</sub> arrest. We could confirm that overexpression of *p27<sup>KIP1</sup>* alone under high-serum conditions was able to block TKO-Bcl2 MEFs in G<sub>2</sub>. This is consistent with recent work showing an inhibitory interaction of *p27<sup>KIP1</sup>* and Cyclin B1 (Nakayama et al., 2004; Martin et al., 2005; Aleem et al., 2005). Furthermore, inhibition of *p21<sup>CIP1</sup>* by RNA interference markedly delayed the G<sub>2</sub> arrest, although it was not sufficient for unrestrained proliferation under mitogen-starved conditions. Others have previously suggested a role for *p21<sup>CIP1</sup>* in a G<sub>2</sub> arrest upon DNA damage by inhibition of Cyclin B1 kinase activity in the nucleus (Smits et al., 2000b; Baus et al., 2003). It is likely that *p21<sup>CIP1</sup>* RNAi-treated cells ultimately arrest in G<sub>2</sub> through inhibition of Cyclin A- and B1-associated kinase activities by *p27<sup>KIP1</sup>*.

Interestingly, we found both *p21<sup>CIP1</sup>* and *p27<sup>KIP1</sup>* associated to Cyclin B1 in cells expressing E1A and Bcl2 or cells expressing nondegradable Cyclin D1, indicating that the G<sub>2</sub> emergency brake also operates in cells with partially inhibited pocket protein activity. Indeed, the interaction between *p27<sup>KIP1</sup>* and Cyclin B1 was also seen in serum-starved *Rb<sup>-/-</sup>* MEFs and became increasingly evident when additional pocket proteins were ablated, providing a possible explanation for the observed synergy of *Rb* and *p27<sup>KIP1</sup>* loss in murine tumorigenesis (Park et al.,



**Figure 8.** *p21<sup>CIP1</sup>* and *p27<sup>KIP1</sup>* as central players in a mitogen starvation-induced G<sub>1</sub> and G<sub>2</sub> cell cycle arrest

1999). Our results are summarized in Figure 8, showing the CKIs as central mediators of restriction points operating in G<sub>1</sub> or G<sub>2</sub>, depending on the presence of functional pocket proteins.

The interaction of Cyclin B1 and *p21<sup>CIP1</sup>* in arrested cells could be disrupted by blocking the ATM/ATR kinases, suggesting that part of the G<sub>2</sub> arrest is elicited by a DNA damage response. However, the Cyclin B1-*p27<sup>KIP1</sup>* interaction appears to be independent from a DNA damage response, since blocking of ATM/ATR kinase activity did not decrease the stability of this complex. Apparently, the *p27<sup>KIP1</sup>*-mediated inhibition of Cyclin B1 (and possibly Cyclin A) is sufficient to maintain the G<sub>2</sub> arrest.

Loss of p53 suppressed apoptosis and abrogated the G<sub>2</sub> arrest in serum-starved TKO MEFs. Thus, besides its role in protecting cells from entering mitosis after DNA damage (Taylor and Stark, 2001), p53 activity is also involved in mitogenic signaling during G<sub>2</sub>. p53 might act as a sensor of mitogenic signaling; however, it might also merely act as a factor that allows cells to accumulate sufficient levels of *p21<sup>CIP1</sup>* during serum starvation. In p53 knockdown TKO MEFs, *p21<sup>CIP1</sup>* does not accumulate because it is a direct transcriptional target of p53. The reduced inhibition of Cdk2 activity in these cells might result in phosphorylation and hence degradation of *p27<sup>KIP1</sup>* (Morisaki et al., 1997). However, why then was inhibition of *p21<sup>CIP1</sup>* by RNAi not sufficient for unhindered proliferation under mitogen-deprived conditions, whereas p53 inhibition was? A trivial explanation is that *p21<sup>CIP1</sup>* was more effectively reduced by p53 RNAi than by *p21<sup>CIP1</sup>* RNAi (compare Figures 3B and 3C and Figures 7G and 7H). Other possibilities we are currently addressing are that p53 targets other than *p21<sup>CIP1</sup>* contribute to the reduction of Cdk2 activity upon serum starvation and that *p27<sup>KIP1</sup>* may be a direct p53 target as was suggested before (Fontemaggi et al., 2002).

Another question is whether TKO-p53RNAi MEFs have become fully serum independent. On the one hand, this seemed to be the case as cells reached confluency by exponential growth in the absence of serum. On the other hand, the generation time had doubled, suggesting that TKO-p53RNAi cells

are still dependent on serum for optimal growth. A procedure allowing passaging of cells under serum-deprived conditions should reveal whether these growth characteristics persist beyond three population doublings.

Loss of the G<sub>1</sub> restriction point is believed to be an essential step in the development of cancer (Hanahan and Weinberg, 2000). Dependent on the cell type or type of cancer, this defect can be the result of overexpression of *Cyclin D1*, loss of one or more of the pocket proteins, or expression of a viral oncogene or combination of these. However, our results suggest that, besides loss of the G<sub>1</sub> restriction point, unconstrained proliferation also requires suppression of apoptosis and alleviation of a G<sub>2</sub> cell cycle arrest. Recent studies on retinoblastoma in mice may provide an example of this scenario. Inactivation of *Rb* in the murine retina resulted in increased proliferation of retinal progenitors, which was enhanced by additional loss of *p107*. While the majority of the retinal cells (rods, cones, and ganglion and bipolar cells) subsequently underwent apoptosis, amacrine precursors arrested. The latter compartment gave rise to retinoblastomas in *Rbp130*- and *Rbp107*-deficient retinas, suggesting that alleviation of cell cycle arrest rather than suppression of apoptosis was required for oncogenic transformation of pocket protein-deficient cells (Dannenberg et al., 2004; Robanus-Maandag et al., 1998; MacPherson et al., 2004). Although no loss of p53 was observed in murine retinoblastoma, we speculate that other events impinging on the G<sub>2</sub> arrest may have occurred to allow oncogenic transformation (Robanus-Maandag et al., 1998; Dannenberg et al., 2004). On the other hand, loss of *Rb* and loss of *p53* do strongly synergize in murine tumorigenesis. For example, combinational loss of *Rb* and *p53* in the CNS of mice resulted in medulloblastoma at 3–4 months of age, whereas individual loss of either *Rb* or *p53* did not cause this malignancy (Marino et al., 2000). Our findings provide a novel explanation for this strong synergy.

## Experimental procedures

### Isolation and culturing of MEFs

MEFs were isolated from chimeric embryos as described before (Dannenberg et al., 2000) and cultured in GMEM (Invitrogen-GIBCO) supplemented with 10% fetal calf serum, 1 mM nonessential amino acids (Invitrogen-GIBCO), 1 mM sodium pyruvate (Invitrogen-GIBCO), 100 μg/ml penicillin, 100 μg/ml streptomycin (Invitrogen-GIBCO), and 0.1 mM β-mercaptoethanol (Merck). For serum starvation experiments, cells were trypsinized and allowed to attach under high-serum conditions for 1 hr. Subsequently, cells were washed with PBS and supplemented with serum-free medium. For serum stimulation experiments, serum-free medium was replaced by complete medium for the indicated times. In the case of caffeine treatment, cells were fed with serum-free medium containing 5 mM caffeine (Sigma). Growth curves were performed as described previously (Dannenberg et al., 2000). For synchronization in mitosis, MEFs were cultured in the presence of 250 ng/ml nocodazole for 8 hr. Mitotic cells were isolated by shake-off and replated at indicated conditions.

### Constructs, transfection, and retroviral infections

The pRetroSuper system and pRetroSuper-p53 were described previously (Dirac and Bernards, 2003). A retroviral vector encoding a nondegradable form of Cyclin D1 was obtained from R. Agami (Agami and Bernards, 2000). The 19-mer p21<sup>CIP1</sup> targeting sequence in pRetroSuper-p21<sup>CIP1</sup> is GCCCTACTCTGTGTGTCT. The LZRS-Bcl2-Zeo plasmid was obtained from A. Werner (Werner et al., 2002). Ecotropic retroviral supernatants were produced by transfecting Phoenix cells by calcium phosphate coprecipitation. Thirty-six hours posttransfection, retroviral supernatants were filtered through a 0.45 μm filter and used to infect MEFs. MEFs were sequentially infected three times, each time for at least 6 hr in the presence of 4 μg/ml

polybrene. In the case of the pRetroSuper constructs, cells were allowed to recover from the retroviral infection for 7 days and subsequently used for BrdU incorporation assays. Bcl2-expressing MEFs were selected in 25 μg/ml bleomycin for 7 days.

### FACS analysis

To measure BrdU incorporation, cells were incubated with 10 mM BrdU for 1 hr. Cells were fixed in 70% ethanol at 4°C. Cells were stained with mouse anti-BrdU (Becton Dickinson) as described in Brugarolas et al. (1998) or stained for MPM2 positivity (Smits et al., 2000a) and subsequently analyzed on FACS using “Cell Quest” software and “FACS Express” software (De Novo Software).

### Time-lapse microscopy and immunofluorescence

For time-lapse microscopy experiments, cells were plated on 35 mm culture dishes (Wilco dishes) (van Vugt et al., 2004). Microinjection was performed as described previously (van Vugt et al., 2004). For immunofluorescence microscopy, cells were grown on coverslips and grown in the absence of serum or irradiated with 10 Gy of γ irradiation. Cells were fixed in 4% formaldehyde and permeabilized using 0.2% Triton X (Sigma) in PBS. Aspecific staining was blocked by incubation in PBS containing 5% BSA for 30 min. Slides were incubated with indicated antibodies in PBS containing 5% of BSA for 2 hr at room temperature. DNA was stained using To-Pro3 Dye (Molecular Probes), and slides were mounted in the presence of Vectashield to prevent bleaching (Vector Laboratories). Slides were analyzed by confocal microscopy as described previously (van Vugt et al., 2004).

### Immunoprecipitations, immunoblots, and antibodies

Cells were harvested and subsequently lysed for 30 min in ELB (150 mM NaCl, 50 mM HEPES [pH 7.5], 5 mM EDTA, 0.1% NP-40) containing protease inhibitors (Complete; Roche) and phosphatase inhibitors (5 mM NaF, 0.5 mM sodium vanadate, and 20 mM β-glycerolphosphate). Protein concentrations were determined using the Bradford assay (Bio-Rad). For immunoprecipitations, 60 mg of protein was incubated with 0.4 μg of immobilized antibody overnight at 4°C while being rotated. In the case of immunoprecipitations for kinase assays, 20 μg (Cyclin B1) or 50 μg (Cyclin A) of protein was used. For kinase assays, immunoprecipitation reactions were washed four times in ELB and once in kinase buffer (50 mM HEPES [pH 7.5], 5 mM MgCl<sub>2</sub>, 2.5 mM MnCl<sub>2</sub>, 1 mM DTT). Each reaction was performed in 30 μl of kinase buffer with 5 mg of Histone H1 (Roche) and [γ-<sup>32</sup>P] dATP for 30 min at 37°C. Subsequently, the reaction mixture was separated by SDS-PAGE, and after gel drying, the signal was quantified using phospho-imaging screens and TINA software. The antibodies used were rabbit anti-Cyclin A (C19), rabbit anti-Cyclin B1 (H433), mouse anti-Cyclin B1 (GNS1), rabbit anti-p21 (C19), mouse anti-p21 (F5), mouse anti-Cdk1 (17), rabbit anti-Cdk2 (M2), rabbit anti-Cdk4 (C22), goat anti-Actin (I19) (Santa Cruz), mouse anti-Cyclin A (Ab2) (Neomarkers), mouse anti-p27 (BD Transduction laboratories), MPM2 (Upstate), Alexa Fluor 488 anti-mouse and 568 anti-rabbit secondary antibodies (Molecular Probes), or HRP-conjugated goat anti-mouse, HRP-conjugated goat anti-rabbit, and HRP-conjugated rabbit anti-goat (Dako) secondary antibodies.

### Supplemental data

The Supplemental Data include one supplemental figure and can be found with this article online at <http://www.cancer.org/cgi/content/full/8/6/455/DC1/>.

### Acknowledgments

We thank A. Dirac for the mouse pRetroSuper and pRetroSuper-p53 RNAi vectors and A. van der Wal for injecting blastocysts for MEF isolation. We thank T. Brummelkamp and M. Voorhoeve for reagents, L. Hengst for providing the YFP-p27<sup>KIP1</sup> construct, and K. Palande for DNA damage experiments in TKO MEFs. We thank R. Kortlever for MEFs expressing a nondegradable form of Cyclin D1 and P. Krimpenfort for p16<sup>INK4a</sup> null MEFs. We are grateful to J. Hansen and J.-H. Dannenberg for fruitful discussions and M. van Lohuizen and C. Wielders for critically reading the manuscript. This work was supported by the Dutch Cancer Society (NKI 2000-2232, NKI 2002-2634).

Received: June 10, 2005  
 Revised: September 8, 2005  
 Accepted: October 31, 2005  
 Published: December 12, 2005

## References

- Agami, R., and Bernards, R. (2000). Distinct initiation and maintenance mechanisms cooperate to induce G1 cell cycle arrest in response to DNA damage. *Cell* 102, 55–66.
- Aleem, E., Kiyokawa, H., and Kaldis, P. (2005). Cdc2-cyclin E complexes regulate the G1/S phase transition. *Nat. Cell Biol.* 7, 831–836.
- Almasan, A., Yin, Y., Kelly, R.E., Lee, E.Y., Bradley, A., Li, W., Bertino, J.R., and Wahl, G.M. (1995). Deficiency of retinoblastoma protein leads to inappropriate S-phase entry, activation of E2F-responsive genes, and apoptosis. *Proc. Natl. Acad. Sci. USA* 92, 5436–5440.
- Baus, F., Gire, V., Fisher, D., Piette, J., and Dulic, V. (2003). Permanent cell cycle exit in G2 phase after DNA damage in normal human fibroblasts. *EMBO J.* 22, 3992–4002.
- Blagosklonny, M.V., and Pardee, A.B. (2002). The restriction point of the cell cycle. *Cell Cycle* 1, 103–110.
- Borel, F., Lohez, O.D., Lacroix, F.B., and Margolis, R.L. (2002). Multiple centrosomes arise from tetraploidy checkpoint failure and mitotic centrosome clusters in p53 and RB pocket protein-compromised cells. *Proc. Natl. Acad. Sci. USA* 99, 9819–9824.
- Brugarolas, J., Bronson, R.T., and Jacks, T. (1998). p21 is a critical CDK2 regulator essential for proliferation control in Rb-deficient cells. *J. Cell Biol.* 147, 503–514.
- Burgering, B.M., and Kops, G.J. (2002). Cell cycle and death control: long live Forkheads. *Trends Biochem. Sci.* 27, 352–360.
- Connor, M.K., Kotchetkov, R., Cariou, S., Resch, A., Lupetti, R., Beniston, R.G., Melchior, F., Hengst, L., and Slingerland, J.M. (2003). CRM1/Ran-mediated nuclear export of p27(Kip1) involves a nuclear export signal and links p27 export and proteolysis. *Mol. Biol. Cell* 14, 201–213.
- Coqueret, O. (2003). New roles for p21 and p27 cell-cycle inhibitors: a function for each cell compartment? *Trends Cell Biol.* 13, 65–70.
- Dannenberger, J.H., van Rossum, A., Schuijff, L., and te Riele, H. (2000). Ablation of the retinoblastoma gene family deregulates G(1) control causing immortalization and increased cell turnover under growth-restricting conditions. *Genes Dev.* 14, 3051–3064.
- Dannenberger, J.H., Schuijff, L., Dekker, M., van der Valk, M., and te Riele, H. (2004). Tissue-specific tumor suppressor activity of retinoblastoma gene homologs p107 and p130. *Genes Dev.* 18, 2952–2962.
- Dirac, A.M., and Bernards, R. (2003). Reversal of senescence in mouse fibroblasts through lentiviral suppression of p53. *J. Biol. Chem.* 278, 11731–11734.
- Fontemaggi, G., Kela, I., Amariglio, N., Rechavi, G., Krishnamurthy, J., Strano, S., Sacchi, A., Givol, D., and Blandino, G. (2002). Identification of direct p73 target genes combining DNA microarray and chromatin immunoprecipitation analyses. *J. Biol. Chem.* 277, 43359–43368.
- Furuno, N., den Elzen, N., and Pines, J. (1999). Human cyclin A is required for mitosis until mid prophase. *J. Cell Biol.* 147, 295–306.
- Hanahan, D., and Weinberg, R.A. (2000). The hallmarks of cancer. *Cell* 100, 57–70.
- Helt, A.M., and Galloway, D.A. (2003). Mechanisms by which DNA tumor virus oncoproteins target the Rb family of pocket proteins. *Carcinogenesis* 24, 159–169.
- Herrera, R.E., Sah, V.P., Williams, B.O., Makela, T.P., Weinberg, R.A., and Jacks, T. (1996). Altered cell cycle kinetics, gene expression, and G1 restriction point regulation in Rb-deficient fibroblasts. *Mol. Cell. Biol.* 16, 2402–2407.
- MacPherson, D., Sage, J., Kim, T., Ho, D., McLaughlin, M.E., and Jacks, T. (2004). Cell type-specific effects of Rb deletion in the murine retina. *Genes Dev.* 18, 1681–1694.
- Malumbres, M., and Barbacid, M. (2001). To cycle or not to cycle: a critical decision in cancer. *Nat. Rev. Cancer* 1, 222–231.
- Marino, S., Vooijs, M., Der Gulden, H., Jonkers, J., and Berns, A. (2000). Induction of medulloblastomas in p53-null mutant mice by somatic inactivation of Rb in the external granular layer cells of the cerebellum. *Genes Dev.* 14, 994–1004.
- Martin, A., Odajima, J., Hunt, S.L., Dubus, P., Ortega, S., Malumbres, M., and Barbacid, M. (2005). Cdk2 is dispensable for cell cycle inhibition and tumor suppression mediated by p27(Kip1) and p21(Cip1). *Cancer Cell* 7, 591–598.
- Medema, R.H., Kops, G.J., Bos, J.L., and Burgering, B.M. (2000). AFX-like Forkhead transcription factors mediate cell-cycle regulation by Ras and PKB through p27kip1. *Nature* 404, 782–787.
- Morisaki, H., Fujimoto, A., Ando, A., Nagata, Y., Ikeda, K., and Nakanishi, M. (1997). Cell cycle-dependent phosphorylation of p27 cyclin-dependent kinase (Cdk) inhibitor by cyclin E/Cdk2. *Biochem. Biophys. Res. Commun.* 240, 386–390.
- Moroni, M.C., Hickman, E.S., Denchi, E.L., Caprara, G., Colli, E., Cecconi, F., Muller, H., and Helin, K. (2001). Apaf-1 is a transcriptional target for E2F and p53. *Nat. Cell Biol.* 3, 552–558.
- Muller, H., Bracken, A.P., Vernell, R., Moroni, M.C., Christians, F., Grassilli, E., Prosperini, E., Vigo, E., Oliner, J.D., and Helin, K. (2001). E2Fs regulate the expression of genes involved in differentiation, development, proliferation, and apoptosis. *Genes Dev.* 15, 267–285.
- Mulligan, G., and Jacks, T. (1998). The retinoblastoma gene family: cousins with overlapping interests. *Trends Genet.* 14, 223–229.
- Mymryk, J.S., Shire, K., and Bayley, S.T. (1994). Induction of apoptosis by adenovirus type 5 E1A in rat cells requires a proliferation block. *Oncogene* 9, 1187–1193.
- Nakayama, K., Nagahama, H., Minamishima, Y.A., Miyake, S., Ishida, N., Hatakeyama, S., Kitagawa, M., Iemura, S., Natsume, T., and Nakayama, K.I. (2004). Skp2-mediated degradation of p27 regulates progression into mitosis. *Dev. Cell* 6, 661–672.
- Nigg, E.A. (2001). Mitotic kinases as regulators of cell division and its checkpoints. *Nat. Rev. Mol. Cell Biol.* 2, 21–32.
- Nurse, P. (1990). Universal control mechanism regulating onset of M-phase. *Nature* 344, 503–508.
- Ortega, S., Prieto, I., Odajima, J., Martin, A., Dubus, P., Sotillo, R., Barbero, J.L., Malumbres, M., and Barbacid, M. (2003). Cyclin-dependent kinase 2 is essential for meiosis but not for mitotic cell division in mice. *Nat. Genet.* 35, 25–31.
- Pardee, A.B. (1974). A restriction point for control of normal animal cell proliferation. *Proc. Natl. Acad. Sci. USA* 71, 1286–1290.
- Park, M.S., Rosai, J., Nguyen, H.T., Capodieci, P., Cordon-Cardo, C., and Koff, A. (1999). p27 and Rb are on overlapping pathways suppressing tumorigenesis in mice. *Proc. Natl. Acad. Sci. USA* 96, 6382–6387.
- Planas-Silva, M.D., and Weinberg, R.A. (1997). The restriction point and control of cell proliferation. *Curr. Opin. Cell Biol.* 9, 768–772.
- Rao, L., Debbas, M., Sabbatini, P., Hockenbery, D., Korsmeyer, S., and White, E. (1992). The adenovirus E1A proteins induce apoptosis, which is inhibited by the E1B 19-kDa and Bcl-2 proteins. *Proc. Natl. Acad. Sci. USA* 89, 7742–7746.
- Robanus-Maandag, E., Dekker, M., van der Valk, M., Carrozza, M.L., Jeanny, J.C., Dannenberg, J.H., Berns, A., and te Riele, H. (1998). p107 is a suppressor of retinoblastoma development in pRb-deficient mice. *Genes Dev.* 12, 1599–1609.
- Sage, J., Mulligan, G.J., Attardi, L.D., Miller, A., Chen, S., Williams, B., Theodorou, E., and Jacks, T. (2000). Targeted disruption of the three Rb-related genes leads to loss of G(1) control and immortalization. *Genes Dev.* 14, 3037–3050.

- Schmidt, M., Fernandez de Mattos, S., van der Horst, A., Klompaker, R., Kops, G.J., Lam, E.W., Burgering, B.M., and Medema, R.H. (2002). Cell cycle inhibition by FoxO forkhead transcription factors involves downregulation of cyclin D. *Mol. Cell. Biol.* 22, 7842–7852.
- Smits, V.A., Klompaker, R., Arnaud, L., Rijksen, G., Nigg, E.A., and Medema, R.H. (2000a). Polo-like kinase-1 is a target of the DNA damage checkpoint. *Nat. Cell Biol.* 2, 672–676.
- Smits, V.A., Klompaker, R., Vallenius, T., Rijksen, G., Makela, T.P., and Medema, R.H. (2000b). p21 inhibits Thr161 phosphorylation of Cdc2 to enforce the G2 DNA damage checkpoint. *J. Biol. Chem.* 275, 30638–30643.
- Stanelle, J., Stiewe, T., Theseling, C.C., Peter, M., and Putzer, B.M. (2002). Gene expression changes in response to E2F1 activation. *Nucleic Acids Res.* 30, 1859–1867.
- Taylor, W.R., and Stark, G.R. (2001). Regulation of the G2/M transition by p53. *Oncogene* 20, 1803–1815.
- Toyoshima, F., Moriguchi, T., Wada, A., Fukuda, M., and Nishida, E. (1998). Nuclear export of cyclin B1 and its possible role in the DNA damage-induced G2 checkpoint. *EMBO J.* 17, 2728–2735.
- van Vugt, M.A., van de Weerd, B.C., Vader, G., Janssen, H., Calafat, J., Klompaker, R., Wolthuis, R.M., and Medema, R.H. (2004). Polo-like kinase-1 is required for bipolar spindle formation but is dispensable for anaphase promoting complex/Cdc20 activation and initiation of cytokinesis. *J. Biol. Chem.* 279, 36841–36854.
- Werner, A.B., de Vries, E., Tait, S.W., Bontjer, I., and Borst, J. (2002). Bcl-2 family member Bfl-1/A1 sequesters truncated bid to inhibit its collaboration with pro-apoptotic Bak or Bax. *J. Biol. Chem.* 277, 22781–22788.