

Opinion

miRNA Nomenclature: A View Incorporating Genetic Origins, Biosynthetic Pathways, and Sequence Variants

T. Desvignes,¹ P. Batzel,¹ E. Berezikov,² K. Eilbeck,^{3,4}
J.T. Eppig,⁵ M.S. McAndrews,⁵ A. Singer,⁶ and
J.H. Postlethwait^{1,*}

High-throughput sequencing of miRNAs has revealed the diversity and variability of mature and functional short noncoding RNAs, including their genomic origins, biogenesis pathways, sequence variability, and newly identified products such as miRNA-offset RNAs (moRs). Here we review known cases of alternative mature miRNA-like RNA fragments and propose a revised definition of miRNAs to encompass this diversity. We then review nomenclature guidelines for miRNAs and propose to extend nomenclature conventions to align with those for protein-coding genes established by international consortia. Finally, we suggest a system to encompass the full complexity of sequence variations (i.e., isomiRs) in the analysis of small RNA sequencing experiments.

Small RNAs and Their Functions

At least three kinds of small RNAs help regulate animal gene expression: Piwi-interacting RNAs (piRNAs) and two RNAi systems, small interfering RNAs (siRNAs) and miRNAs [1–4]. The animal-specific piRNA system involves RNA molecules about 24–30 nucleotides (nt) long that form RNA–protein complexes by interacting with Piwi proteins to regulate epigenetic and post-transcriptional gene silencing of retrotransposons and other genetic elements primarily in germ cells, especially during spermatogenesis [5]. Second, the siRNA pathway regulates genetic expression using about 21-nt-long RNA molecules of either exogenous (often viral) or endogenous origin [6]. siRNAs can interact with an Argonaute protein and incorporate into an RNA–protein complex [RNA-Induced Silencing Complex (RISC)] to alter chromatin structure, transcriptional regulation, and transposable element repression; siRNAs can also participate in post-transcriptional regulation by inducing mRNA degradation based on complementarity between the siRNA and specific mRNAs. siRNAs can act as an antiviral defense system that targets viral mRNAs using exogenous fragments of the target viral mRNA or can act as a self-cell regulator of translation by the production of endogenous siRNA [6]. Finally, the miRNA pathway uses endogenous small RNA molecules about 22 nt long to modulate the translation of targeted mRNAs by acting on their stability or translation through interaction with the RISC [6].

miRNAs help control cell-fate determination, cell differentiation, and organ development and physiology and are involved in pathology and disease in humans, other metazoans, and plants [7–11]. Recent deep-sequencing experiments and powerful bioinformatics tools [12] have led to a dramatic increase in the number of known miRNA genes (*miRNA* genes), even in well-studied

Trends

Functional miRNAs can originate from various types of noncoding RNA genes.

The definition of miRNAs should take a functional perspective.

The full complexity of isomiR populations should be considered in miRNA studies.

miRNA nomenclature systems should encompass origins, function, and isomiRs.

¹Institute of Neuroscience, University of Oregon, Eugene, OR 97403, USA

²European Research Institute for the Biology of Ageing, University of Groningen, University Medical Center Groningen, 9713 AV Groningen, The Netherlands

³Utah Science, Technology, and Research Center for Genetic Discovery, University of Utah, Salt Lake City, UT 84112, USA

⁴Department of Biomedical Informatics, University of Utah, Salt Lake City, UT 84112, USA

⁵Mouse Genome Informatics, The Jackson Laboratory, 600 Main Street, Bar Harbor, ME 04609, USA

⁶ZFIN, 5291 University of Oregon, Eugene, OR 97403-5291, USA

*Correspondence: jpostle@uoneuro.uoregon.edu (J.H. Postlethwait).

organisms such as mouse and human. Databases such as miRBase [13], ZooMir [14], miRNEST [15], and miRMaid [16] have cataloged miRNA sequences across various organisms. miRBase, the most commonly used reference miRNA database, currently lists 1915 mouse and 2588 human mature miRNA sequences (miRNA), but dramatically fewer in most other species; for example, miRBase currently lists only 350 mature miRNAs in zebrafish (release 21; [13]). The extent to which the disparity in annotated miRNAs among species reflects biology with evolved differences in genome content or, instead, simply variation in sequencing effort or in stringency of gene annotation (which seems more likely for closely related species) is an issue requiring investigation.

The constantly increasing number of reads delivered by massively parallel sequencing strategies reveals putative new *miRNA* genes, some of which are indeed novel but others of which are orthologous to *miRNA* genes previously described in other species. Curiously, many short reads containing some features of mature miRNAs appear to originate from other classes of RNA such as small nucleolar RNAs (snoRNAs) or long noncoding RNAs (lncRNAs) [17]. Deep sequencing has also revealed isomiRs [18], which are variant miRNA sequences that all appear to derive from the same gene but vary in sequence due to post-transcriptional processing. Because isomiRs contain different sequences, they may have different targets and thus different cell functions [19]. Great sequencing depth and careful data mining also reveal moRs [20–22] and single-strand RNA fragments released from the loop RNA portion of the precursor miRNA hairpin [23,24] [which we call ‘loop-origin miRNAs’ (loRs)]. Although functional evidence remains scarce, these variants appear to be independently regulated and may play cooperative, complementary, or alternative roles in cell function and gene regulation compared with their associated miRNAs. This complexity in genomic origin and variation in sequence may impact cell biology but is so far under-studied; furthermore, current nomenclature conventions make annotation rather subjective depending on the emphasis of each researcher in accepting or rejecting putative new miRNA-related sequences.

A problem regarding short RNAs derived from non-miRNA genes is that transcription may be quite widespread across the genome and that many such perhaps randomly transcribed sequences may be without function, although they might be detected with deep sequencing. In addition, it is possible that some promiscuously transcribed sequences might be processed by Drosha/Dicer, and with ever-deeper sequencing identifying this ‘noise’ becomes more likely. Consistent lengths, 5′ processing sites, species conservation, and tissue-specific expression patterns provide evidence for control and hence circumstantial evidence for function. A prudent course of action would seem to be to annotate apparent isomiRs and/or miRNA-like fragments derived from non-miRNA genes to facilitate testing for functional roles of individual sequences.

The complexity of miRNA origins and sequence variations inhibit optimal communication and connectivity among species. Here we explore recent findings and opinions concerning miRNA variations and raise the question: what should define an animal miRNA, a single biogenesis pathway or a specific mode of action? To answer this question, we propose to define miRNAs from a functional rather than a biogenesis perspective. In addition, we suggest nomenclature guidelines that parallel international protein-coding gene nomenclature guidelines to improve connectivity in miRNA research across metazoans.

Canonical miRNAs: Products and Biogenesis

Most *miRNA* genes are usually transcribed by RNA polymerase II into a long transcript called the ‘primary miRNA transcript’ (*pri-MiRNA*) [17,19]. For *miRNA* genes organized in clusters, the *pri-MiRNA* can be monocistronic or polycistronic. Within the nucleus, the DROSHA and DGCR8 (known as Pasha in *Drosophila melanogaster* and *Caenorhabditis elegans*) microprocessor

complex cleaves the *pri-MiRNA* into a shorter sequence called the 'miRNA precursor' (*pre-MiRNA*), which displays a hairpin-like secondary structure [25]. The *pre-MiRNA* is exported to the cytoplasm by Exportin-5 in a complex with RAN-GTP and then DICER1, an RNase III enzyme (hereafter called Dicer), cleaves off the loop of the *pre-MiRNA* and generates a mature miRNA duplex. The two strands of the miRNA duplex pair with a few mismatches and each strand usually has a 2-nt overhang at its 3' end. The duplex then loads onto an Argonaute protein, the effector of the RISC [25], where the unwinding of the duplex and the selection of a strand occur [26,27]. The mature miRNA, as part of the RISC, binds the 3' untranslated region (UTR) of specific mRNAs and mediates mRNA degradation, destabilization, or translational inhibition according to sequence complementarity to the target [19,28,29]. This general biogenesis pathway and the features of the intermediate transcripts define a so-called 'canonical *Mir* gene' (Figure 1). Criteria for the annotation of canonical miRNA genes were first defined in 2003 [7] and rely on combined expression and biogenesis criteria (Box 1). Criteria have been revised over the years (Box 1) to fit with new knowledge often arising from technological improvements [13,30–33].

Alternative Sources of miR-Like Small RNAs

Although canonical miRNA products encompass the largest fraction of known mature miRNAs, several miRNA-like products originate by DROSHA-independent and/or Dicer-independent biogenesis pathways and from the processing of other noncoding RNA biotypes and associated with Argonaute proteins [34]. Examples include miRtrons, Argonaute RISC catalytic subunit 2 (Ago2) processing, and origins from various other noncoding RNA genes including snoRNA, lincRNA, and tRNA genes (Figure 1).

miRtrons


The study of *Drosophila* mutants identified miRtrons, miRNAs that are produced from an intron of a protein-coding gene by Droscha-independent pre-mRNA splicing machinery [35–37]. Expression of miRtrons depends on the host gene because at least one end of the *pre-MiRNA* hairpin aligns exactly to the pre-mRNA splice site junction; in contrast, other intronic miRNAs can be regulated independently of their host gene [38–40]. Because miRtrons are expressed with the host protein-coding gene and are processed by the mRNA splicing machinery rather than by the microprocessor complex, they are noncanonical miRNAs, although they follow other miRNA annotation criteria (Box 1).

DROSHA-Dependent, Dicer-Independent miRNAs

Another alternative mode of miRNA biogenesis came from the study of *Dicer1* null mutant mice, which revealed miRNAs that are processed by DROSHA but not Dicer [41,42]. For example, in mouse and zebrafish, *pre-MiR451a* is transcribed, along with its neighbor *pre-MiR144*, into a polycistronic *pri-MiRNA* that is processed by DROSHA, but its subsequent maturation into a unique strand, the 5p strand, occurs not by Dicer but by AGO2, the only vertebrate Argonaute protein displaying catalytic activity [42–44].

miRNA-Like Small RNAs From Other Noncoding RNAs

Alternative miRNA origins have been identified due to recent high-throughput sequencing, which identified 'miRNA-like' small RNAs that are loaded into the RISC [34] and mediate mRNA translation but derive from various noncoding RNA types, including snoRNA [45–48], lincRNA [49], short hairpin RNA (shRNA) [50], or tRNA [51]. For example, mouse *MiR664* originates from *Snora36b*; mouse *MiR320* is initially expressed as a shRNA [50]; mouse *MiR1983* derives from the processing of an alternative secondary conformation of a tRNA(Ile) [50]; and mouse *MiR205* comes from the lincRNA *4631405K08Rik* [52] (although note that, by definition, all canonical pri-miRNAs are lincRNAs). Several sequences originating from such noncoding RNAs, like *MiR1274a*, which is a fragment of a tRNA(Lys), have been removed from recent miRBase



Trends in Genetics

Figure 1. Diverse Genetic Origins and Biogenesis Pathways Leading to Functional miRNAs. Many different types of noncoding RNA genes [e.g., canonical miRNAs, small nucleolar RNAs (snoRNAs), long noncoding RNAs (lncRNAs), tRNAs] are transcribed, cleaved into precursor miRNAs (*pre-miRNAs*) and eventually processed into miRNA duplexes from which one strand loads into an Argonaute protein (AGO) and represses mRNA translation through the RNA-Induced Silencing Complex (RISC). Canonical miRNAs, lncRNA, and AGO2-processed miRNAs (i.e., *MIR451*) depend on DROSHA and DGCR8 for processing of the primary miRNA transcript into the precursor miRNA. miRtron precursors, by contrast, are processed by the mRNA splicing machinery into simple miRtrons, in which intron splicing defines both ends of the hairpin, or tailed miRtrons, which contain templated regions 5' or 3' to a splice-derived terminal intron. snoRNA, short hairpin RNA (shRNA), inverted-repeat-derived endogenous small interfering RNA (endo-siRNA), vaultRNA, and Y_RNA appear to be DROSHA independent and are transcribed directly into an RNA fragment displaying the characteristics of a precursor miRNA. In some cases, tRNAs can also be cleaved to produce small RNA fragments by folding into an alternative secondary structure that shows characteristics of a precursor miRNA. Dicer processes most precursor miRNAs (i.e., canonical miRNA, miRtrons, some snoRNAs, lncRNA, tRNA, shRNA, inverted-repeat-derived endo-siRNA, and vaultRNA) to give rise to a miRNA duplex. Some snoRNA-derived miRNAs and Y_RNA-derived short fragments appear to be Dicer independent but the enzyme producing the short fragments remains unknown. *Cis*- and *trans*-derived endo-siRNAs are by nature transcribed directly into a miRNA duplex. miRNA duplexes are then unwound and one side of the duplex, the mature miRNA, is loaded into an AGO protein that drives the assembly of the RISC and the repression of protein translation of the targeted transcript. An AGO2-processed miRNA is an exception to this biogenesis step because it is Dicer independent and the catalytic activity of the AGO2 protein produces the mature miRNA. Functional experiments are critical to separate true miRNAs from degradation products of these other RNA types.

releases (present in release 16, but gone from release 17 and later), while others, like *MIR1983*, remain as valid entries (release 21). Short 'miRNA-like' RNAs that derive from various noncoding RNAs but load into the RISC and act like canonical miRNAs probably arise from a pre-miRNA-like hairpin alternative secondary structure of the precursor that Dicer processes to form a short fragment that subsequently loads into the RISC [34,50,53,54]. Although these sequences have a different genomic origin than canonical miRNAs, they show other key features of functional miRNAs, including origin from a hairpin-like precursor, processing by Dicer, evolutionary conservation, loading into the RISC, and functional repression of

Box 1. A Timeline for the Evolution of miRNA Annotation

Criteria for annotating *miRNA* genes and miRNA transcripts have evolved since miRNAs were first discovered.

2003: An understanding of miRNA biology begins to emerge. miRNA identity requires a combination of at least one expression and one biogenesis criterion [109]:

A. Expression criteria:

A.1 Hybridization to a size-fractionated RNA sample detects an RNA transcript of about 22 nt.

A.2 A cDNA library made from size-fractionated RNA identifies a sequence of about 22 nt.

B. Biogenesis criteria:

B.1 The precursor sequence is predicted to fold into a hairpin-like structure with the lowest free energy and display the miRNA in one arm of the hairpin.

B.2 An approximately 22-nt miRNA sequence and its predicted fold-back precursor is conserved across phylogenies.

B.3 The precursor accumulates in mutants with reduced Dicer function.

• Special case: candidates identified by sequence analysis can be annotated without further experimental evidence if they are homologous/orthologous to miRNAs from related organisms.

2011: Results begin to appear from deep-sequencing technologies. Criteria revised to avoid mis-annotation of small RNA fragments originating from various RNA species by focusing on miRNA biogenesis [13]:

A. Multiple reads support the sequence of the mature miRNA in several independent libraries.

B. Reads support mature sequences from both arms of the predicted hairpin and the putative mature sequences pair with correct 2-nt 3' overhangs.

C. Reads map to genomic DNA and sequence flanking the putative mature miRNA is predicted to form a miRNA precursor-like hairpin with strong pairing between the mature miRNA and the opposite arm.

D. Reads mapping 'very many times' to a genome are discarded.

E. Mapped reads do not overlap other annotated transcripts (mRNAs or other known RNA types).

F. Reads support a single 5' end of the mature sequence, reflecting precise nucleotide processing, which contrasts with the heterogeneous ends that characterize degraded RNAs and processing of other RNA types.

2013: Increasing evidence suggests the presence of dubious miRNAs annotated in miRNA databases; criteria evolve toward defining 'high-confidence mirs' [33]:

A. At least ten reads must map with no mismatches to each of the two possible mature miRNAs.

B. The two mature miRNAs must pair in a duplex with 0–4-nt overhangs at their 3' ends.

C. At least 50% of the reads for each arm of the pre-miRNA must have the same 5' end.

D. The predicted hairpin structure must have a predicted folding free energy lower than -0.2 kcal/mol/nt.

E. At least 60% of the bases in the mature miRNA should pair in the predicted hairpin structure.

2015: Special cases of miRNA-like sequences from noncanonical biogenesis pathways, like DROSHA-independent miRNAs (e.g., miRtrons), Dicer-independent miRNAs (e.g., *Mir451*), and pathways from other noncoding RNA genes (e.g., snoRNA, lncRNA) could be accepted as miRNAs.

translation. These considerations call into question the current definitions of *miRNA* genes and mature miRNA products and the practice of rejecting miRNAs that emerge from non-miRNA noncoding RNA genes while emphasizing the need to distinguish *bone fide* miRNAs from degradation products.

endo-siRNAs

Some endogenous siRNAs (endo-siRNAs) bind AGO2, load into the RISC, and repress translation [55]. Any of three different mechanisms can make endo-siRNAs: (i) transcription of inverted repeats producing a hairpin-derived endo-siRNA that Dicer processes into an RNA duplex; (ii) transcription of short fragments from both DNA strands of a genomic locus giving a *cis*-derived endo-siRNA that forms a double-stranded RNA duplex; and (iii) transcription of two different genomic loci producing RNA fragments that hybridize to assemble into a *trans*-derived endo-siRNA double-stranded RNA duplex [50,56]. At least some of these endo-siRNAs appear to have a function comparable with canonical miRNAs by cleaving mRNA targets when pairing is perfect [55]. endo-siRNAs, however, also play a role in chromatin regulation, silencing transposable elements in both germ and somatic cell lines, and can act as cotranscriptional silencers of gene expression [57]. Although endo-siRNAs do not form by the canonical miRNA biosynthetic pathway, when loaded onto AGO2 and part of the RISC they appear to function like miRNAs by inducing the decay of mRNA targets; the miRNA regulation system of animals and

plants is likely to have emerged from the siRNA machinery [58]. The finding that siRNAs and the miRNAs of both plants and cnidarians (the basally diverging sister group to bilateral animals) use near-perfect pairing to their targets and act by mRNA cleavage suggests an independent origin of the miRNA systems of plants and animals and the evolution of new miRNA mechanisms in the Bilateria [59]. Given the likely evolutionary origin of the miRNA system of gene regulation, it seems likely that some endo-siRNAs could be confused with miRNAs, especially in basal metazoans.

Y_RNA and vaultRNA

Small RNAs derived from Y_RNA or vaultRNA genes can be found in small RNA sequencing and/or AGO pull-down sequencing datasets [34]. Interestingly, most of those miRNA-like reads map to genes orthologous to human or mouse Y_RNA or vaultRNA genes that also produce small RNA products from a similar position of the transcript, suggesting that they are evolutionarily conserved short products of those noncoding RNAs and not simply degradation fragments. For example, we identified reads from the mouse Y_RNA gene *Rny3* (orthologous to human *RNY3* [60]) or from the vaultRNA gene *Vaultrc5* (co-orthologous to human *VTRNA1* genes [61]) in small RNA sequencing and/or AGO pull-down sequencing datasets such as SRP010168 from [22] or datasets deposited at FaceBase (<https://www.facebase.org/project/155/datasets>). Sequences originating from orthologous regions of Y_RNA or vaultRNA genes were previously annotated in human as *MIR1979* and *MIR886*, respectively, but have now been removed from the miRBase database [33]. Short reads originating from several Y_RNA and vaultRNA genes were also recently reported in zebrafish [62]; one, *mir733*, from the *vault.4* gene and another, *mir735*, from the *Y_RNA.2* gene continue to appear in miRBase (MI0004778 and MI0004781, respectively). Although those fragments appear in various datasets and show evolutionary conservation, a function in repressing translation has been shown only for vault fragments [63] and not yet for Y_RNA fragments [64,65], although the tests performed on Y_RNA fragments might have involved suboptimal conditions [66]. More work is needed to decipher the putative function in translation repression of these evolutionarily conserved miRNA-like fragments and hence determine whether they should be considered miRNAs.

Clarifying the Question: What Is a miRNA?

Evidence reviewed in the previous section shows that many types of small noncoding RNA fragments can sometimes function in a manner similar to canonical miRNAs; that is, they are small, they can load into the RISC, and they can act to mediate cleavage or to inhibit translation of targeted mRNAs although they originate from other noncoding RNA biotypes and do not strictly follow the canonical miRNA biogenesis pathway. This situation raises the question of whether miRNAs should be defined: (i) by mature sequence characteristics and their putative function; or (ii) by their biosynthetic pathway. The biogenesis definition [31,32] would allow only canonical miRNAs and possibly miRtrons to be annotated as miRNAs. The mature sequence characteristics definition emphasizes the structure, sequence conservation across species and samples, regulated expression, and putative action of the mature product, including short, single-stranded RNAs, effective Argonaute protein loading, and translation repression (Box 2).

We propose to distinguish miRNAs from other classes of RNA by dissociating the three notions of 'miRNA gene' (*Mir*), 'miRNA precursor' (*pre-MiRNA*), and 'mature miRNA product' (MiRNA). The proposed new definition recognizes that *miRNA* genes produce miRNAs but that some miRNAs may originate from genes transcribed into other types of noncoding RNAs. This revised definition distinguishes function from biogenesis and simplifies the nomenclature of miRNAs and miRNA-like short sequences (Box 2). These new relationships and connections are now implemented in the Sequence Ontology Database [67], which provides an ontology framework for human and mouse nomenclature consortia [68,69].

Box 2. Expanding miRNA Nomenclature to Encompass Alternative miRNA Origins

Many small noncoding RNA sequences are processed from hairpin-like precursors or RNA duplexes (like *pre-miRNAs*), are functionally loaded into an Argonaute protein in the RISC (as are miRNAs), and are developmentally regulated, despite the fact that they are derived – noncanonically – from non-*Mir*, noncoding RNA genes (such as snoRNA or lincRNA genes) (Figure 1). Many of those sequences were first annotated as miRNAs because of their function or conservation across samples and species, but some were subsequently removed from databases after their genomic origin was understood. A paradigm shift, however, that takes a functional rather than a biogenesis approach to those sequences suggests that some should remain as valid entries in miRNA databases or in RNA-Induced Silencing Small RNA databases. To apprehend the full complexity and diversity of RNA-induced silencing mechanisms and to accommodate miRNA-like sequences emerging from non-miRNA genes, we propose three levels of miRNA annotation (Figure 1).

1. The ‘functional sequence’ level, which corresponds to the ‘mature sequence’ level already implemented in miRBase and referenced with a ‘MIMAT’ accession number. For a sequence to qualify, it must be an endogenous single-stranded RNA molecule ~22 nt long with a significant number of reads allowing statistical analysis and mapping to a ‘limited number’ of locations (maybe about ten) in the reference genome, preferentially be found in several independent libraries, and display a consistent 5’ end of the mature sequence (to discriminate products originating from degradation or other modes of noncoding RNA processing).
2. The ‘precursor miRNA’ level, which corresponds to the miRBase ‘stem-loop’ level referenced with an ‘MI’ accession number in miRBase. The sequence should be able to fold into a hairpin-like structure allowing its processing into a mature miRNA. If transcriptomic evidence exists, any identified moRs should be included in the precursor sequence because this additional sequence will influence the stability of secondary structure folding.
3. A new ‘gene’ level that denotes the genomic origin and encompasses miRNAs coming from canonical *miRNA* genes, miRtrons, snoRNA, lincRNA, tRNA, and others.

These revised criteria and additional hierarchical levels would accept new mature miRNAs if they originate from noncanonical *miRNA* genes (abandoning the criterion that they have to come exclusively from *miRNA* genes) and recognize miRNAs cleaved by AGO2 or originating from Dicer-independent pathways (abandoning the previous criterion accepting only Dicer-processed sequences). Revised criteria would continue to accept sequence orthology among closely related species as sufficient evidence for annotation. Special attention to sequence functionality should, however, be focused on novel miRNA-like sequences never previously described in any species. In addition, for species that can be readily studied in the laboratory, validation of dependence on RNase III, Argonaute loading, and/or functional activity in translation and mRNA stability would reinforce the miRNA annotation.

miRNA Nomenclature Evolution

New knowledge of miRNA origins, biogenesis, and function requires a revision of nomenclature guidelines to provide an unbiased naming protocol, consistent with coding-gene guidelines, that maximizes the connectivity of miRNA analyses across species (Table 1). International genome-feature nomenclature guidelines for model animals are already in use (e.g., <http://www.genenames.org/> [69], <http://www.informatics.jax.org/>, <http://rgd.mcw.edu/> [68,70], <http://zfin.org/> [71]) but have not been applied consistently by researchers and repositories regarding miRNA nomenclature. Examples given here use human, mouse, and zebrafish guidelines. For other species, however, guidelines established for the most closely related model species should be used.

General Case

Nomenclature guidelines should do the following.

- (i) Respect species-specific gene nomenclature conventions. For example, *MIR125a* and *Mir125a* represent orthologous miRNA genes in human and mouse, respectively, and *mir125aa* and *mir125ab* symbolize the two zebrafish co-orthologs of the human *MIR125a* gene that were derived from the teleost genome duplication [72,73].
- (ii) Retain the convention that *MIR*, *Mir*, *mir* (human, mouse, zebrafish) italicized and with a lower case ‘r’ (except for human) refers to the gene or genomic locus; ‘*MIR*’ (*MiR*, *miR*) italicized and with an upper case ‘R’ refers to intermediate transcripts; and ‘MIR’ (MIR, MiR) not italicized and with an upper case ‘R’ refers to the final products: the mature miRNAs.
- (iii) Remove references to species in the official gene name. For example, ‘*mmu-*’ should be removed from the previous designator for the mouse gene ‘*mmu-Mir-125a*’. Reference to the organism should be clearly stated within the text to avoid confusion. Databases that cover multiple species might need to retain species designators.

Table 1. Summary of Nomenclature Guidelines for miRNA Genes and their Various Products in Human, Mouse, and Zebrafish

Symbol ^a			Signifies
Human	Mouse	Zebrafish	
<i>MIRX</i>	<i>MirX</i>	<i>mirX</i>	miRNA gene X
<i>MIRXy</i>	<i>MirXy</i>	<i>mirXy</i>	yth member of the mirX family (y can be a letter or a dot number)
<i>MIRXOS</i>	<i>MirXos</i>	<i>mirXos</i>	Mirror-miRNA gene of the miRNA gene X
<i>pri-MIRX</i>	<i>pri-MirX</i>	<i>pri-mirX</i>	Primary miRNA transcript X
<i>pre-MIRX</i>	<i>pre-MirX</i>	<i>pre-mirX</i>	miRNA precursor X
<i>MIRX-5p</i>	<i>MIRX-5p</i>	<i>MIRX-5p</i>	5' arm mature miRNA sequence
<i>MIRX-3p</i>	<i>MIRX-3p</i>	<i>MIRX-3p</i>	3' arm mature miRNA sequence
<i>MORX-5p</i>	<i>MORX-5p</i>	<i>MoRX-5p</i>	5' arm mature moR sequence
<i>MORX-3p</i>	<i>MORX-3p</i>	<i>MoRX-3p</i>	3' arm mature moR sequence
<i>LORX</i>	<i>LORX</i>	<i>LoRX</i>	Loop fragment of the miRNA precursor
<i>MIRCX</i>	<i>MircX</i>	<i>mircX</i>	mir gene cluster X

^aIn species lacking nomenclature conventions, symbols should follow nomenclature guidelines established by the most closely related reference model organism consortium as agreed by researchers studying the species.

- (iv) Remove hyphens separating '*Mir*' from the number designator. Hyphens are not generally used in human, rodent, and zebrafish gene designations. For example, the gene '*hsa-MIR-125a*' becomes '*MIR125A*', '*mmu-Mir-125a*' becomes '*Mir125a*', and '*dre-mir-125aa*' becomes '*mir125aa*'.
- (v) Specify various forms by unambiguous designators. Genomic locus, primary miRNA transcript, miRNA precursor, and mature forms should use, for example, for the mouse gene, '*Mir125a*'; the primary miRNA transcript, '*pri-Mir125a*'; the miRNA precursor, '*pre-Mir125a*'; and the mature form, '*MIR125a*'. Hyphenation of the primary miRNA and the miRNA precursor (i.e., *pri-MIR* and *pre-MIR*, respectively) are retained in accordance with the common usage of the 'pre-mRNA' symbol to refer to a precursor to an mRNA and the italicization follows nomenclature guidelines for the italicization of intermediate transcripts.
- (vi) Annotation should inform without bias. Formerly, one of the two mature products of a *Mir* gene was called the 'mature strand' or 'guide strand' and was considered to be *the* miRNA because it was more highly expressed, while the opposite strand was called the 'passenger strand' or 'star strand' (miR*) because it was less expressed and thought to be inactive [74]. Alternative strands, however, can be differentially expressed in different tissues, developmental stages, pathological states, or species, requiring inversion of the miR/miR* annotation to reflect expression variations if the miR* strand is functionally active [75–78]. For example, *MIR140-3p* is the most highly expressed miRNA strand in mouse ovaries and testes but *MIR140-5p* is the most highly expressed strand in cartilage [79–81]. Appropriately, miRBase has, since release 17, replaced the 'miR/miR*' symbolism with the unbiased '5p/3p' strand annotation to indicate the position of the strand in the *pre-MIRNA* hairpin independently of its expression status in any given dataset [33].

Special Cases

Mirror-miRNA genes provide another nomenclature problem. Mirror-miRNAs are *pri-MIRNA*s independently transcribed from both DNA strands [34,62,82,83]. Although mirror-miRNA pairs are evolutionarily associated, they may be transcribed independently and are unlikely to have a conserved seed; thus, mirror-miRNAs may act on different target protein-coding genes. The convention for protein-coding genes is to use the suffix 'os', for 'opposite strand', for a pair of overlapping genes transcribed from opposite strands. We propose to apply this convention to

mirror-miRs: the first *Mir* gene described would be assigned a number and a transcript documented to come from the opposite strand in the same genomic location would get the 'os' suffix. For example, *Mir3120*, the mirror gene of *Mir214* [83], should be called *Mir214os* and its respective mature sequences MIR214os-5p and MIR214os-3p (Table 1).

miRNA genes with identical or similar mature products can be confused. Because these genes come from gene duplication events, including genome duplication, segmental duplication, or tandem duplication, using the same miRNA number has the advantage of grouping these genes in a common gene family that reflects related structure and evolutionary origin. Adding an additional letter or number as a suffix should follow established species-specific conventions for coding genes. For example, in zebrafish, 'a' and 'b' designators reflect an origin by the teleost genome duplication and thus an ohnological relationship. Examples include *mir125aa* and *mir125ab*, which reside on zebrafish chromosomes LG16 and LG19, which are ohnologous chromosomes from the teleost genome duplication [72], and *MIR125A* and *MIR125B1* in human chromosome bands 19q13.41 and 11q24.1, respectively, derived from the two rounds of early vertebrate genome duplications [84,85]. In accord with tandemly duplicated protein-coding genes, the use of a 'dot-number' suffix should indicate tandem *mir* gene duplicates; for example, the zebrafish tandem duplicates *mir34b* and *mir34c* should be named *mir34b.1* and *mir34b.2*, respectively, according to zebrafish nomenclature conventions. In mouse and human, the approved guideline is the use of letter designators to reflect duplication (Table 1).

Clustered *miRNA* genes, such as the group of six (*Mir17*, *Mir18*, *Mir19a*, *Mir20a*, *Mir19b-1*, and *Mir92a-1*) in the distal region of mouse chromosome 14, are defined by a possible common expression as a unique polycistronic *pri-MiRNA*. No clear guidelines have yet been established to refer to miRNA clusters; sometimes they have been referred to by the first and last genes of the cluster, in the sense of transcription, separated by a tilde (~) or a dash (-) when all genes are transcribed in the same orientation; in this example, the cluster would be annotated *Mir17~92* or *Mir17-92* [86]. For database purposes, however, the use of a tilde is not appropriate because it usually has a predefined meaning. We suggest the use of the nomenclature symbol 'Mirc', standing for 'miRNA cluster', which is already in use in the Mouse Nomenclature System [68]. miRNA clusters can be named and referred to by following nomenclature conventions already in use for specific species using sequentially numbered clusters [57], as, for example, the cluster above is called *Mirc1* in mouse. Alternatively, we suggest that, in other species, the number of the cluster refers to the first gene of the cluster in the sense of transcription, as is currently used for human miRNA clusters [58]; the *MIR17-92* cluster in the example above, which is currently called *MIR17HG* in humans (where HG stands for host gene, although in many cases each *MIR* gene in the cluster would be its own transcriptional unit, questioning the use of the term 'host gene'), would be called *MIRC17* in humans and *mirc17* in zebrafish (Table 1). It is noteworthy that, in some species or duplicated clusters within a species, the first gene of the cluster, which would give the cluster its name, can be lost. It would then be the responsibility of the researcher to establish orthology/paralogy relationships between clusters and give the cluster a name that preserves evolutionary relationships even if the first gene (or other genes in the cluster) is lost; for example, if in zebrafish the 'b' copy of this cluster is missing *mir17*, the cluster would nevertheless be called '*mirc17b*' by analogy with *Hox* clusters, which retain the cluster designation across organisms even when the exact gene content of each cluster may differ in different species.

Other Relevant miRNA Products

isomiRs

High-throughput sequencing has revealed various mature sequences apparently transcribed from a single *Mir* gene. These *bona fide* mature miRNA variants, distinguished from sequencing errors by their frequency, are called isomiRs and can vary in length or contain untemplated nucleotide additions or post-transcriptional base edits [18,22]. Most commonly, variations

involve length at the 3' end. Variations at the 3' end (3'-isomiRs) can result in shorter sequences presumably due to exonuclease 'nibbling' at the mature sequence or in longer sequences probably due to inconsistent Dicer processing or post-transcriptional addition of a few bases that may or may not match the genomic locus, resulting in templated or untemplated variants, respectively. 5'-isomiRs, which vary in length at the 5' terminus, are rarer but can still account for a significant proportion of the isomiR pool. Finally, post-transcriptionally edited isomiRs harbor different internal nucleotide sequences [87].

Although the broad biological significance of isomiRs remains to be fully understood, increasing evidence suggests that some may be functionally important because they can be expressed in a cell-specific manner and hence are regulated [88–94]. Studies also show that some isomiRs vary in stability, loading efficiency onto Argonaute proteins, and target selection [87,95]. Because miRNA repression of mRNA function is hypothesized to rely mainly on 'seed pairing' (the perfect or near-perfect complementary match of nucleotides 2–8 of the mature miRNA product [96–98]), length modifications at the 5' end of the mature product could relocate the seed position and thus change in the seed sequence (called 'seed shifting'), thereby altering mRNA target recognition and function [92,93]. Furthermore, because noncanonical base pairing [28,29,99–102] can cause any portion of the miRNA to influence the pairing of the miRNA to its mRNA-UTR, any modification or addition in the mature sequence can putatively affect biological function.

These considerations raise the question: how should we annotate isomiRs that vary in length or nucleotide sequence? This question is important because tissue- or time-specific variation in the relative abundance of different isomiRs could contribute to differences in cell differentiation, proliferation, or physiology among tissues within a species or to the evolution of different features between species.

Currently, miRNA databases typically display only one mature sequence per strand, which is generally inferred from small RNA sequencing data and/or from calculated miRNA precursor folding probabilities. Although some databases list sequences of isomiRs, this information is not easily accessible and is often associated with only a *pre-MiRNA* sequence and not a mature miRNA sequence, meaning that an isomiR cannot be specifically assigned to any mature product of the precursor and cannot be directly annotated and thus conveniently designated in an investigation. Furthermore, no consistent nomenclature conventions for isomiRs have yet been established, which complicates communication. In databases, the annotated mature sequence usually corresponds to the most highly expressed isomiR across all deposited sequences. The sequencing of more tissues or more species could make the most highly expressed sequence a moving target, which does not facilitate precise communication. To overcome this problem, we propose a new view of miRNA sequences organized into three interconnected levels: (i) the reference mature miRNA sequence, which would be an unchanging standard for reference purposes; (ii) the isomiR sequences, which include all of the miRNA variants observed for a reference miRNA with the application of an accession number to each isomiR and its assignment to one or multiple reference miRNA sequences, if appropriate; and (iii) the functional mature sequence, which would correspond to the most highly expressed isomiR in a given situation, tissue, and developmental stage and thus would provide a functional perspective (Box 3 and Figure 1).

moRs and loRs

Increased sequencing depth has identified other *Mir* gene-encoded fragments in addition to the 5' and 3' miRs. For example, moRs are encoded by sequences adjacent to the ends of the 5' and 3' miRNA-encoding sequences that abut the loop [103]. moRs were first considered to be nonfunctional fragments or processing byproducts, but because the quantity of a moR relative

Box 3. Introducing a Reference Sequence and isomiR Annotation

Current small RNA databases list one 'mature sequence' per miRNA strand, which is usually the most frequent sequence in the database, while sometimes displaying isomiR sequences in a visual quantitative presentation in which each isomiR can be followed to an individualized page with a sequence-specific accession number. Several factors, however, could help better capture the complexity of isomiRs because, so far, those isomiRs are associated with a stem-loop only and not a mature sequence (Figure 1).

1. A 'reference sequence miRNA' (RefSeq-miRNA) should replace the concept of a single universal 'mature sequence'. The effective mature sequence would now be seen in a functional perspective and could vary in different tissues or developmental stages, whereas the reference sequence would be a species-specific invariable standard for each miRNA-encoding gene strand. This convention has the advantage that even if the ranking of isomiR expression levels varies in different tissues, stages, or situations and thus the mature sequence varies, the reference sequence remains unchanged and isomiR sequences and quantities can be described compared with an unchanging reference.
2. Similar to mature sequences that can originate from one or multiple stem-loops when they are closely related, isomiRs can potentially originate from one or multiple stem-loops and thus be isomiRs of one or multiple mature sequences. Each isomiR should thus be associated with all of the mature sequences of which it is possible to be a variant, as for mature sequences.
3. Finally, to facilitate communication, each isomiR should continue to receive its own accession number, like the 'MR' annotation system currently used in miRBase, and corresponding sequences and annotations should be easily accessible for download.

to its neighboring miRNA can vary between tissues or developmental stages, they appear to be regulated and thus are either functional or represent tissue-specific, regulated transcriptional noise [20,21,104–106]. Genomically, moRs are generally more frequent on the 5' side of the hairpin, independent of strand-specific miRNA expression levels [21], consistent with a putative function but also suggesting that they might be coproducts of miRNA biogenesis [107] and thus provide a distinct functional class of miRNA-related sequences. moRs are frequently found in RISC pull-down libraries, suggesting function, and a study has verified that some viral moRs show moderate but still significant repression of mRNA translation in *luciferase* assays [108]. Together, data suggest that moRs can program the RISC and act on gene expression and should be considered in small RNA studies.

moRs were initially annotated as a miRNA with a dot-number suffix referring to its relative position in the hairpin; for example, the moR for the *Mir125a* gene was labeled MiR125a.1 and the original miRNA was called MiR125a.2 because it is further from the end of the hairpin. The symbol moR, introduced by Shi *et al.* [20], however, coupled with designating origin from the 5' or 3' end, for example 'MOR125a-5p' and 'MOR125a-3p', has the advantage that it allows clear and unambiguous identification of the corresponding RNA sequence (Table 1).

Small RNA sequencing libraries also identify sequences that derive from the loop of *pre-MiRNAs* and, like moRs, their relative quantities can vary according to tissue or developmental stage [23,24], suggesting independent regulation of expression and thus independent cell functions. Furthermore, recent studies reported the expression and incorporation of loop-derived short sequences into the RISC and repression of translation of targeted mRNA in *in vitro* luciferase assays [23,24]. We propose use of the nomenclature 'loR' for loop-origin miRNAs; for example, the loop-origin fragment from the *Mir125a* gene would become 'LOR125a' (Table 1).

Concluding Remarks

The increasing amount of small RNA sequencing data provides a more complete view of the putative regulatory small RNAome, but it is important to distinguish the functional miRNAome from the degradome: small RNA fragments resulting from RNA degradation. miRNAs account for most of the reads in small RNA sequencing efforts, but many other miRNA-like small RNAs that originate from noncanonical *miRNA* genes show consistent and significant expression across experiments and can act on message stability and translation repression as do canonical miRNAs. These miRNA-like sequences thus constitute a new kind of miRNA for which we suggest nomenclature conventions. Furthermore, the depth of small RNA sequencing data

Outstanding Questions

What fraction of small noncoding miRNA-like sequences that derive from genes other than traditional miRNA genes act on translational control in a fashion similar to most miRNAs? Although these miRNA-like fragments consistently appear in small RNA sequencing and Argonaute pull-down experiments, evidence of a functional role in translational control remains scarce or absent. Are those short noncoding RNAs incorporated into the RISC and do they inhibit translation of targeted messenger RNAs? Are they evolutionarily conserved? Are they merely degradation products of other noncoding RNA biotypes without function? Do they modulate the miRNA pathway by overwhelming the RISC and inhibiting the loading of *bona fide* miRNAs? The latter possibility would add an additional layer of complexity to the pathway of miRNA regulation.

What are the functions of moRs and loRs, which arise from the ends of pri-miRNA stems or the loop? Evolutionary conservation, sequence conservation, and variable expression levels suggest a functional role in mRNA translation control, but conclusive evidence in direct protein translation is lacking or minimal. Do moRs and loRs function like miRNA by entering the RISC and regulating protein translation? Are they simple byproducts of the miRNA biogenesis pathway? If so, are they needed for proper pri-miRNA and pre-miRNA processing, thus explaining their evolutionary conservation? Answering these questions would provide a deeper understanding of the miRNA biogenesis pathway and the roles of miRNA genes in regulatory biology.

demonstrates that a single *Mir* gene can potentially produce not just a single mature miRNA from each of its two complementary paired stem sequences, but up to five different and transcriptionally regulated, putatively functional mature sequences in the order moR-5p, miR-5p, loR, miR-3p, and moR-3p. Finally, most of these sequences show variations (isomiRs) arising from both templated and untemplated variations. Appreciation of the variety of sequences that can arise from genes that encode miRNAs should help to unravel the regulatory roles they play in development and physiology, health, and disease. The revised nomenclature conventions we suggest should help facilitate communication about the variety of miRNA-related sequences derived from different species and, hence, the conservation evolution of miRNA functions (see Outstanding Questions).

Acknowledgments

The authors thank Eric C. Lai for valuable comments. The work was funded by the grants NIH R01HG004341 (K.E.), NIH 5P41HG000330 (J.T.E. and M.S.M.), NIH HG004838 (A.S.), and NIH HG004834 (A.S.), the Research Council of Norway (FishmiR project, grant no. 213825, J.H.P. and I. Babiak), NIH U01DE020076 (J.H.P., K. Artinger, and D. Clouthier), NIH 5R240D011199 (J.H.P. and W. Cresko), and NIH R01 OD011116 (J.H.P.).

References

- Lee, R.C. *et al.* (1993) The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 75, 843–854
- He, L. and Hannon, G.J. (2004) MicroRNAs: small RNAs with a big role in gene regulation. *Nat. Rev. Genet.* 5, 522–531
- Ghildiyal, M. and Zamore, P.D. (2009) Small silencing RNAs: an expanding universe. *Nat. Rev. Genet.* 10, 94–108
- Ketting, R.F. (2011) The many faces of RNAi. *Dev. Cell* 20, 148–161
- Ishizu, H. *et al.* (2012) Biology of PIWI-interacting RNAs: new insights into biogenesis and function inside and outside of germ-lines. *Genes Dev.* 26, 2361–2373
- Carthew, R.W. and Sontheimer, E.J. (2009) Origins and mechanisms of miRNAs and siRNAs. *Cell* 136, 642–655
- Ambros, V. (2004) The functions of animal microRNAs. *Nature* 431, 350–355
- Kosik, K.S. (2010) MicroRNAs and cellular phenotypy. *Cell* 143, 21–26
- Bizuayehu, T.T. and Babiak, I. (2014) MicroRNA in teleost fish. *Genome Biol. Evol.* 6, 1911–1937
- Christodoulou, F. *et al.* (2010) Ancient animal microRNAs and the evolution of tissue identity. *Nature* 463, 1084–1088
- Grimson, A. *et al.* (2008) Early origins and evolution of microRNAs and Piwi-interacting RNAs in animals. *Nature* 455, 1193–1197
- Gomes, C.P.C. *et al.* (2013) A review of computational tools in microRNA discovery. *Front. Bioinform. Comput. Biol.* 4, 81
- Kozomara, A. and Griffiths-Jones, S. (2011) miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Res.* 39, D152–D157
- Li, S.-C. *et al.* (2010) Identification of homologous microRNAs in 56 animal genomes. *Genomics* 96, 1–9
- Szczesniak, M.W. and Makalowska, I. (2014) miRNEST 2.0: a database of plant and animal microRNAs. *Nucleic Acids Res.* 42, D74–D77
- Jacobsen, A. *et al.* (2010) miRmaid: a unified programming interface for microRNA data resources. *BMC Bioinformatics* 11, 29
- Ha, M. and Kim, V.N. (2014) Regulation of microRNA biogenesis. *Nat. Rev. Mol. Cell Biol.* 15, 509–524
- Morin, R.D. *et al.* (2008) Application of massively parallel sequencing to microRNA profiling and discovery in human embryonic stem cells. *Genome Res.* 18, 610–621
- Ameres, S.L. and Zamore, P.D. (2013) Diversifying microRNA sequence and function. *Nat. Rev. Mol. Cell Biol.* 14, 475–488
- Shi, W. *et al.* (2009) A distinct class of small RNAs arises from pre-miRNA-proximal regions in a simple chordate. *Nat. Struct. Mol. Biol.* 16, 183–189
- Langenberger, D. *et al.* (2009) Evidence for human microRNA-offset RNAs in small RNA sequencing data. *Bioinformatics* 25, 2298–2301
- Zhou, H. *et al.* (2012) Deep annotation of mouse iso-miR and iso-moR variation. *Nucleic Acids Res.* 40, 5864–5875
- Okamura, K. *et al.* (2013) Functional small RNAs are generated from select miRNA hairpin loops in flies and mammals. *Genes Dev.* 27, 778–792
- Winter, J. *et al.* (2013) Loop-miRs: active microRNAs generated from single-stranded loop regions. *Nucleic Acids Res.* 41, 5503–5512
- Winter, J. *et al.* (2009) Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nat. Cell Biol.* 11, 228–234
- Kawamata, T. *et al.* (2009) Structural determinants of miRNAs for RISC loading and slicer-independent unwinding. *Nat. Struct. Mol. Biol.* 16, 953–960
- Kwak, P.B. and Tomari, Y. (2012) The N domain of Argonaute drives duplex unwinding during RISC assembly. *Nat. Struct. Mol. Biol.* 19, 145–151
- Bartel, D.P. (2009) MicroRNAs: target recognition and regulatory functions. *Cell* 136, 215–233
- Shin, C. *et al.* (2010) Expanding the microRNA targeting code: functional sites with centered pairing. *Mol. Cell* 38, 789–802
- Brown, M. *et al.* (2013) Mammalian miRNA curation through next-generation sequencing. *Front. Genet.* 4, 145
- Hansen, T.B. *et al.* (2011) Enhancing miRNA annotation confidence in miRBase by continuous cross dataset analysis. *RNA Biol.* 8, 378–383
- Langenberger, D. *et al.* (2011) MicroRNA or not microRNA? In *Advances in Bioinformatics and Computational Biology* (de Souza, O.N. *et al.*, eds), pp. 1–9, Springer
- Kozomara, A. and Griffiths-Jones, S. (2013) miRBase: annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res.* 42, D68–D73
- Burroughs, A.M. *et al.* (2011) Deep-sequencing of human Argonaute-associated small RNAs provides insight into miRNA sorting and reveals Argonaute association with RNA fragments of diverse origin. *RNA Biol.* 8, 158–177
- Okamura, K. *et al.* (2007) The miRtron pathway generates microRNA-class regulatory RNAs in *Drosophila*. *Cell* 130, 89–100
- Ruby, J.G. *et al.* (2007) Intronic microRNA precursors that bypass Drosha processing. *Nature* 448, 83–86
- Berezikov, E. *et al.* (2007) Mammalian miRtron genes. *Mol. Cell* 28, 328–336
- Isik, M. *et al.* (2010) Expression patterns of intronic microRNAs in *Caenorhabditis elegans*. *Silence* 1, 5

39. Ramalingam, P. *et al.* (2013) Biogenesis of intronic miRNAs located in clusters by independent transcription and alternative splicing. *RNA* 20, 76–87
40. Desvignes, T. *et al.* (2014) Evolution of the miR199-214 cluster and vertebrate skeletal development. *RNA Biol.* 11, 281–294
41. Cheloufi, S. *et al.* (2010) A dicer-independent miRNA biogenesis pathway that requires Ago catalysis. *Nature* 465, 584–589
42. Cifuentes, D. *et al.* (2010) A novel miRNA processing pathway independent of Dicer requires Argonaute2 catalytic activity. *Science* 328, 1694–1698
43. Liu, J. *et al.* (2004) Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 305, 1437–1441
44. Yang, J.-S. *et al.* (2010) Conserved vertebrate mir-451 provides a platform for Dicer-independent, Ago2-mediated microRNA biogenesis. *Proc. Natl. Acad. Sci. U.S.A.* 107, 15163–15168
45. Ender, C. *et al.* (2008) A human snoRNA with microRNA-like functions. *Mol. Cell* 32, 519–528
46. Taft, R.J. *et al.* (2009) Small RNAs derived from snoRNAs. *RNA* 15, 1233–1240
47. Brameier, M. *et al.* (2011) Human box C/D snoRNAs with miRNA like functions: expanding the range of regulatory RNAs. *Nucleic Acids Res.* 39, 675–686
48. Scott, M.S. and Ono, M. (2011) From snoRNA to miRNA: dual function regulatory non-coding RNAs. *Biochimie* 93, 1987–1992
49. Keniry, A. *et al.* (2012) The H19 lincRNA is a developmental reservoir of miR-675 that suppresses growth and Igf1r. *Nat. Cell Biol.* 14, 659–665
50. Babiarz, J.E. *et al.* (2008) Mouse ES cells express endogenous shRNAs, siRNAs, and other Microprocessor-independent, Dicer-dependent small RNAs. *Genes Dev.* 22, 2773–2785
51. Xia, J. *et al.* (2013) Noncanonical microRNAs and endogenous siRNAs in normal and psoriatic human skin. *Hum. Mol. Genet.* 22, 737–748
52. Wang, D. *et al.* (2013) MicroRNA-205 controls neonatal expansion of skin stem cells by modulating the PI₃K pathway. *Nat. Cell Biol.* 15, 1153–1163
53. Miyoshi, K. *et al.* (2010) Many ways to generate microRNA-like small RNAs: non-canonical pathways for microRNA production. *Mol. Genet. Genomics* 284, 95–103
54. Aalto, A.P. and Pasquinelli, A.E. (2012) Small non-coding RNAs mount a silent revolution in gene expression. *Curr. Opin. Cell Biol.* 24, 333–340
55. Okamura, K. and Lai, E.C. (2008) Endogenous small interfering RNAs in animals. *Nat. Rev. Mol. Cell Biol.* 9, 673–678
56. Kim, V.N. *et al.* (2009) Biogenesis of small RNAs in animals. *Nat. Rev. Mol. Cell Biol.* 10, 126–139
57. Chen, L. *et al.* (2012) Naturally occurring endo-siRNA silences LINE-1 retrotransposons in human cells through DNA methylation. *Epigenetics* 7, 758–771
58. Axtell, M.J. *et al.* (2011) *Vive la différence*: biogenesis and evolution of microRNAs in plants and animals. *Genome Biol.* 12, 221
59. Moran, Y. *et al.* (2014) Cnidarian microRNAs frequently regulate targets by cleavage. *Genome Res.* Published online March 18, 2014. <http://dx.doi.org/10.1101/gr.162503.113>
60. Perreault, J. *et al.* (2007) Ro-associated Y RNAs in metazoans: evolution and diversification. *Mol. Biol. Evol.* 24, 1678–1689
61. Stadler, P.F. *et al.* (2009) Evolution of vault RNAs. *Mol. Biol. Evol.* 26, 1975–1991
62. Desvignes, T. *et al.* (2014) Expanding the annotation of zebrafish microRNAs based on small RNA sequencing. *Gene* 546, 386–389
63. Persson, H. *et al.* (2009) The non-coding RNA of the multidrug resistance-linked vault particle encodes multiple regulatory small RNAs. *Nat. Cell Biol.* 11, 1268–1271
64. Meiri, E. *et al.* (2010) Discovery of microRNAs and other small RNAs in solid tumors. *Nucleic Acids Res.* 38, 6234–6246
65. Nicolas, F.E. *et al.* (2012) Biogenesis of Y RNA-derived small RNAs is independent of the microRNA pathway. *FEBS Lett.* 586, 1226–1230
66. Verhagen, A.P.M. and Pruijn, G.J.M. (2011) Are the Ro RNP-associated Y RNAs concealing microRNAs? Y RNA-derived miRNAs may be involved in autoimmunity. *Bioessays* 33, 674–682
67. Eilbeck, K. *et al.* (2005) The Sequence Ontology: a tool for the unification of genome annotations. *Genome Biol.* 6, R44
68. Bult, C.J. *et al.* (2012) The Mouse Genome Database: genotypes, phenotypes, and models of human disease. *Nucleic Acids Res.* 41, D885–D891
69. Gray, K.A. *et al.* (2013) Genenames.org: the HGNC resources in 2013. *Nucleic Acids Res.* 41, D545–D552
70. Laulederkind, S.J.F. *et al.* (2013) The Rat Genome Database 2013 – data, tools and users. *Brief. Bioinform.* 14, 520–526
71. Bradford, Y. *et al.* (2011) ZFIN: enhancements and updates to the zebrafish model organism database. *Nucleic Acids Res.* 39, D822–D829
72. Amores, A. *et al.* (1998) Zebrafish *hox* clusters and vertebrate genome evolution. *Science* 282, 1711–1714
73. Braasch, I. and Postlethwait, J.H. (2012) Polyploidy in fish and the teleost genome duplication. In *Polyploidy and Genome Evolution* (Soltis, P.S. and Soltis, D.E., eds), pp. 341–383, Springer
74. Griffiths-Jones, S. (2004) The microRNA Registry. *Nucleic Acids Res.* 32, D109–D111
75. Okamura, K. *et al.* (2008) The regulatory activity of microRNA* species has substantial influence on microRNA and 3' UTR evolution. *Nat. Struct. Mol. Biol.* 15, 354–363
76. Czech, B. *et al.* (2009) Hierarchical rules for Argonaute loading in *Drosophila*. *Mol. Cell* 36, 445–456
77. Okamura, K. *et al.* (2009) Distinct mechanisms for microRNA strand selection by *Drosophila* Argonautes. *Mol. Cell* 36, 431–444
78. Yang, J.-S. *et al.* (2011) Widespread regulatory activity of vertebrate microRNA* species. *RNA* 17, 312–326
79. Tuddenham, L. *et al.* (2006) The cartilage specific microRNA-140 targets histone deacetylase 4 in mouse cells. *FEBS Lett.* 580, 4214–4217
80. Eberhart, J.K. *et al.* (2008) MicroRNA Mirn140 modulates Pdgf signaling during palatogenesis. *Nat. Genet.* 40, 290–298
81. Rakoczy, J. *et al.* (2013) MicroRNAs-140-5p/140-3p modulate Leydig cell numbers in the developing mouse testis. *Biol. Reprod.* 88, 143
82. Tyler, D.M. *et al.* (2008) Functionally distinct regulatory RNAs generated by bidirectional transcription and processing of microRNA loci. *Genes Dev.* 22, 26–36
83. Scott, H. *et al.* (2012) MiR-3120 is a mirror microRNA that targets heat shock cognate protein 70 and auxilin messenger RNAs and regulates clathrin vesicle uncoating. *J. Biol. Chem.* 287, 14726–14733
84. Dehal, P. and Boore, J.L. (2005) Two rounds of whole genome duplication in the ancestral vertebrate. *PLoS Biol.* 3, e314
85. Nakatani, Y. *et al.* (2007) Reconstruction of the vertebrate ancestral genome reveals dynamic genome reorganization in early vertebrates. *Genome Res.* 17, 1254–1265
86. Saini, H.K. *et al.* (2008) Annotation of mammalian primary microRNAs. *BMC Genomics* 9, 564
87. Neilsen, C.T. *et al.* (2012) IsomiRs – the overlooked repertoire in the dynamic microRNAome. *Trends Genet.* 28, 544–549
88. Lee, L.W. *et al.* (2010) Complexity of the microRNA repertoire revealed by next-generation sequencing. *RNA* 16, 2170–2180
89. Marti, E. *et al.* (2010) A myriad of miRNA variants in control and Huntington's disease brain regions detected by massively parallel sequencing. *Nucleic Acids Res.* 38, 7219–7235
90. Llorens, F. *et al.* (2013) A highly expressed miR-101 isomiR is a functional silencing small RNA. *BMC Genomics* 14, 104
91. Guo, L. and Chen, F. (2014) A challenge for miRNA: multiple isomiRs in miRNAomics. *Gene* 544, 1–7
92. Tan, G.C. *et al.* (2014) 5' isomiR variation is of functional and evolutionary importance. *Nucleic Acids Res.* 42, 9424–9435
93. Xia, J. and Zhang, W. (2014) A meta-analysis revealed insights into the sources, conservation and impact of microRNA 5'-isoforms in four model species. *Nucleic Acids Res.* 42, 1427–1441
94. Warnefors, M. *et al.* (2014) Conserved microRNA editing in mammalian evolution, development and disease. *Genome Biol.* 15, R83

95. Thornton, J.E. *et al.* (2014) Selective microRNA uridylation by Zcchc6 (TUT7) and Zcchc11 (TUT4). *Nucleic Acids Res.* 42, 11777–11791
96. Brennecke, J. *et al.* (2005) Principles of microRNA-target recognition. *PLoS Biol.* 3, e85
97. Grimson, A. *et al.* (2007) MicroRNA targeting specificity in mammals: determinants beyond seed pairing. *Mol. Cell* 27, 91–105
98. Schirle, N.T. *et al.* (2014) Structural basis for microRNA targeting. *Science* 346, 608–613
99. Li, N. *et al.* (2008) Dispatched Homolog 2 is targeted by miR-214 through a combination of three weak microRNA recognition sites. *Nucleic Acids Res.* 36, 4277–4285
100. Thomas, M. *et al.* (2010) Desperately seeking microRNA targets. *Nat. Struct. Mol. Biol.* 17, 1169–1174
101. Chi, S.W. *et al.* (2012) An alternative mode of microRNA target recognition. *Nat. Struct. Mol. Biol.* 19, 321–327
102. Clark, P.M. *et al.* (2014) Argonaute CLIP-seq reveals miRNA targetome diversity across tissue types. *Sci. Rep.* 4, 5947
103. Bortoluzzi, S. *et al.* (2011) MicroRNA-offset RNAs (moRNAs): by-product spectators or functional players? *Trends Mol. Med.* 17, 473–474
104. Bortoluzzi, S. *et al.* (2012) Characterization and discovery of novel miRNAs and moRNAs in JAK2V617F-mutated SET2 cells. *Blood* 119, e120–e130
105. Asikainen, S. *et al.* (2015) Selective microRNA-offset RNA expression in human embryonic stem cells. *PLoS ONE* 10, e0116668
106. Zhang, W. *et al.* (2010) Multiple distinct small RNAs originate from the same microRNA precursors. *Genome Biol.* 11, R81
107. Berezikov, E. *et al.* (2011) Deep annotation of *Drosophila melanogaster* microRNAs yields insights into their processing, modification, and emergence. *Genome Res.* 21, 203–215
108. Umbach, J.L. *et al.* (2010) Analysis of rhesus rhadinovirus microRNAs expressed in virus-induced tumors from infected rhesus macaques. *Virology* 405, 592–599
109. Ambros, V. *et al.* (2003) A uniform system for microRNA annotation. *RNA* 9, 277–279